

t2\_square\_1  
(TMEgQUISC8d8imVaEneZrsyg3kUx29XtX8fE)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k4\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 k6\_numbers X0) \Rightarrow (r1\_xxreal\_0 X1 (k2\_xcmplx\_0 X0 X1)))) \quad (1)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow ((r1\_xxreal\_0 X1 X0) \Rightarrow (k4\_xxreal\_0 X0 X1 = X0)) \wedge ((\neg r1\_xxreal\_0 X1 X0) \Rightarrow (k4\_xxreal\_0 X0 X1 = X1)))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcmplx\_0 X0) \wedge (v1\_xcmplx\_0 X1)) \Rightarrow (k2\_xcmplx\_0 X0 X1 = k2\_xcmplx\_0 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xxreal\_0 X0) \quad (5)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (6)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \wedge (r1\_xxreal\_0 k6\_numbers X1)) \Rightarrow (r1\_xxreal\_0 (k4\_xxreal\_0 X0 X1) (k2\_xcmplx\_0 X0 X1))))$$