

t2_topmetr2

(TMUutgK3bSoCCN5pZEktDaW77jC4sq5Xetq)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X2 X1)) \Rightarrow (r1_tarski (k2_xboole_0 X0 X2) X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k2_xboole_0 X0 (k4_xboole_0 X1 X0) = k2_xboole_0 X0 X1 \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (r1_tarski (k10_xtuple_0 X0) (k10_xtuple_0 (k1_funct_4 X1 X0)))) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (r1_tarski (k10_xtuple_0 (k1_funct_4 X0 X1)) (k2_xboole_0 (k10_xtuple_0 X0) (k10_xtuple_0 X1)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (5)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski (k7_relat_1 X0) (k3_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1))) (k10_xtuple_0 X1)) \Rightarrow (r1_tarski (k6_subset_1 (k10_xtuple_0 X0) (k10_xtuple_0 X1)) (k10_xtuple_0 (k1_funct_4 X0 X1)))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(X0 = X1) \Leftrightarrow ((r1_tarSKI X0 X1) \wedge (r1_tarSKI X1 X0)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \quad (8)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarSKI (k7_relat_1 X0 (\\ k3_xboole_0 (k9_xtuple_0 X0) (k9_xtuple_0 X1))) (k10_xtuple_0 \\ X1)) \Rightarrow (k2_xboole_0 (k10_xtuple_0 X0) (k10_xtuple_0 X1) = k10_xtuple_0 \\ (k1_funct_4 X0 X1)))) \end{aligned}$$