

t2_xreal_1
(TMcgqX5Uh5aejc3sD5NERGan9953Gqj9Ep7)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\neg \forall X1.(v1_xreal_0 X1) \Rightarrow (r1_xxreal_0 X1 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 (k4_xcmplx_0 X1) (k4_xcmplx_0 X0)))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k4_xcmplx_0 (k4_xcmplx_0 X0) = X0) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((v1_xcmplx_0 (k4_xcmplx_0 X0)) \wedge (v1_xreal_0 (k4_xcmplx_0 X0))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (5)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\neg \forall X1.(v1_xreal_0 X1) \Rightarrow (r1_xxreal_0 X0 X1))$$