

t2_yellow17

(TMY3y3iThiqY5vU3W3Y9rpfybGL5zFqkWZE)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_card_3 : \iota \Rightarrow \iota$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((X0 \in k4_card_3 X1) \Leftrightarrow ((k9_xtuple_0 X0 = k9_xtuple_0 X1) \wedge (\forall X2.(X2 \in k9_xtuple_0 X1) \Rightarrow (k1_funct_1 X0 X2 \in k1_funct_1 X1 X2)))))) \quad (1)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \forall X3.(X2 \neq X3) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X3 = k1_funct_1 X0 X3)) \quad (2)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. (X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X2 = X1)) \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. k9_xtuple_0 (k2_funct_7 X0 X2 X1) = k9_xtuple_0 X0) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v1_relat_1 (k2_funct_7 X0 X1 X2)) \wedge (v1_funct_1 (k2_funct_7 X0 X1 X2))) \quad (5)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \forall X3.((X3 \in k1_funct_1 X0 X2) \wedge (X1 \in k4_card_3 X0)) \Rightarrow (k2_funct_7 X1 X2 X3 \in k4_card_3 X0))))$$