

t2_zfrefle1

(TMJbv93Chm8x2gejmkwgd1B6UZnKXbJDZUp)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k9_zflang : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_zfrefle1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_zflang : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $r2_zf_model : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ k9_zflang)) \Rightarrow ((r1_zfrefle1 X0 X1) \Leftrightarrow (\forall X2. ((v1_zflang \\ X2) \wedge (m2_finseq_1 X2 k5_numbers)) \Rightarrow ((X2 \in X1) \Rightarrow (r2_zf_model X0 X2)))))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ k9_zflang)) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 k9_zflang)) \Rightarrow \\ (((r1_tarski X1 X2) \wedge (r1_zfrefle1 X0 X2)) \Rightarrow (r1_zfrefle1 X0 X1)))) \end{aligned}$$