

t30_analort

(TMZxzn1kyNAq9pfvE4H4DKWMWadXAM3ye3o)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_analmetr : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_analort : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_analoaf : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\
 & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\
 & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\
 & X0)))))))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow \\
 & (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 \\
 & X3 (u1_struct_0 X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (u1_struct_0 \\
 & X0)) \Rightarrow (\forall X5.(m1_subset_1 X5 (u1_struct_0 X0)) \Rightarrow (\forall X6. \\
 & (m1_subset_1 X6 (u1_struct_0 X0)) \Rightarrow ((r1_analmetr X0 X1 X2) \Rightarrow ((r1_analoaf \\
 & X0 X3 X4 X5 X6) \Leftrightarrow (\exists X7.(m1_subset_1 X7 (u1_struct_0 X0)) \wedge (\\
 & \exists X8.(m1_subset_1 X8 (u1_struct_0 X0)) \wedge ((X7 \neq X8) \wedge ((r1_analort \\
 & X0 X1 X2 X7 X8 X3 X4) \wedge (r1_analort X0 X1 X2 X7 X8 X5 X6))))))))))))) \\
 & \tag{1}
 \end{aligned}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\
 & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\
 & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\
 & X0)))))))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow \\
 & (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow ((r1_analoaf X0 \\
 & X1 X2 X2 X1) \Rightarrow (X1 = X2)))) \\
 & \tag{2}
 \end{aligned}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\ & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\ & X0)))))))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 \\ & X3 (u1_struct_0 X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (u1_struct_0 \\ & X0)) \Rightarrow (\forall X5.(m1_subset_1 X5 (u1_struct_0 X0)) \Rightarrow (\forall X6. \\ & (m1_subset_1 X6 (u1_struct_0 X0)) \Rightarrow (\neg(r1_analmetr X0 X1 X2) \wedge (\\ r1_analort X0 X1 X2 X3 X4 X5 X6) \wedge ((r1_analort X0 X1 X2 X3 X4 X6 X5) \wedge (\\ & (X3 \neq X4) \wedge (X5 \neq X6))))))))))))) \end{aligned}$$