

# t30\_bhsp\_2 (TMLgGHeQHvX- htXJNV3vWFzjRipPVnUypxrS)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v2\_bhsp\_1 : \iota \Rightarrow o$  be given. Let  $l1\_bhsp\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_bhsp\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_bhsp\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_comseq\_2 : \iota \Rightarrow o$  be given. Let  $k2\_bhsp\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_normsp\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_normsp\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_rlvect\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_seq\_2 : \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned}
 & \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\
 & X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\
 & ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge ((v2\_bhsp\_1 \\
 & X0) \wedge (l1\_bhsp\_1 X0)))))))))) \Rightarrow (\forall X1. (m1\_subset\_1 X1 k1\_numbers) \Rightarrow \\
 & (\forall X2. ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 k5\_numbers (u1\_struct\_0 \\
 & X0)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (u1\_struct\_0 \\
 & X0)))))) \Rightarrow ((v1\_bhsp\_2 X2 X0) \Rightarrow (v1\_bhsp\_2 (k5\_normsp\_1 X0 X2 X1) \\
 & X0))))))
 \end{aligned}
 \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\
& X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\
& ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge ((v2\_bhs\_p1 \\
& X0) \wedge (l1\_bhs\_p1 X0)))))))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 \\
& X0)) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 k5\_numbers \\
& (u1\_struct\_0 X0)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& k5\_numbers (u1\_struct\_0 X0)))))) \Rightarrow (((v1\_bhs\_p2 X2 X0) \wedge (k1\_bhs\_p2 \\
& X0 X2 = X1)) \Rightarrow ((v2\_comseq\_2 (k2\_bhs\_p2 X0 (k4\_normsp\_1 X0 X2 X1)) \wedge \\
& (k2\_seq\_2 (k2\_bhs\_p2 X0 (k4\_normsp\_1 X0 X2 X1)) = k6\_numbers))))))
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\
& X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\
& ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge ((v2\_bhs\_p1 \\
& X0) \wedge (l1\_bhs\_p1 X0)))))))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 k1\_numbers) \Rightarrow \\
& (\forall X2.((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 X2 k5\_numbers (u1\_struct\_0 \\
& X0)) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (u1\_struct\_0 \\
& X0)))))) \Rightarrow ((v1\_bhs\_p2 X2 X0) \Rightarrow (k1\_bhs\_p2 X0 (k5\_normsp\_1 X0 X2 X1) = \\
& k1\_rlvect\_1 X0 (k1\_bhs\_p2 X0 X2 X1))))))
\end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.(l1\_bhs\_p1 X0) \Rightarrow (l1\_rlvect\_1 X0) \tag{4}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.\forall X2.(((\neg v2\_struct\_0 X0) \wedge (l1\_rlvect\_1 \\
& X0)) \wedge (((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 X1 k5\_numbers (u1\_struct\_0 \\
& X0)) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (u1\_struct\_0 \\
& X0)))))) \wedge (m1\_subset\_1 X2 k1\_numbers)) \Rightarrow ((v1\_funct\_1 (k5\_normsp\_1 \\
& X0 X1 X2)) \wedge ((v1\_funct\_2 (k5\_normsp\_1 X0 X1 X2) k5\_numbers (u1\_struct\_0 \\
& X0)) \wedge (m1\_subset\_1 (k5\_normsp\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& k5\_numbers (u1\_struct\_0 X0))))))
\end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.(((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge \\
& ((v2\_rlvect\_1 X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 \\
& X0) \wedge ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge \\
& ((v2\_bhs\_p1 X0) \wedge (l1\_bhs\_p1 X0)))))))))) \wedge ((v1\_funct\_1 X1) \wedge \\
& ((v1\_funct\_2 X1 k5\_numbers (u1\_struct\_0 X0)) \wedge (m1\_subset\_1 X1 \\
& (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (u1\_struct\_0 X0)))))) \Rightarrow \\
& (m1\_subset\_1 (k1\_bhs\_p2 X0 X1) (u1\_struct\_0 X0))
\end{aligned} \tag{6}$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\ & X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\ & ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge ((v2\_bhspl-1 \\ & X0) \wedge (l1\_bhspl-1 X0)))))))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 \\ & X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 k1\_numbers) \Rightarrow (\forall X3.(( \\ & v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 k5\_numbers (u1\_struct\_0 X0)) \wedge \\ & (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers (u1\_struct\_0 \\ & X0)))))) \Rightarrow (((v1\_bhspl-2 X3 X0) \wedge (k1\_bhspl-2 X0 X3 = X1)) \Rightarrow ((v2\_comseq\_2 \\ & (k2\_bhspl-2 X0 (k4\_normsp\_1 X0 (k5\_normsp\_1 X0 X3 X2) (k1\_rlvect\_1 \\ & X0 X1 X2)))) \wedge (k2\_seq\_2 (k2\_bhspl-2 X0 (k4\_normsp\_1 X0 (k5\_normsp\_1 \\ & X0 X3 X2) (k1\_rlvect\_1 X0 X1 X2))) = k6\_numbers)))))) \end{aligned}$$