

t30_borsuk_5

(TMdmZ9eiZCiKXXTmhLBStsoXbsV3486i43v)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k3_borsuk_5 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_rat_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_borsuk_5 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((X0 \in k4_xxreal_1 X1 X2) \Leftrightarrow ((\neg r1_xxreal_0 X0 X1) \wedge \\ & (\neg r1_xxreal_0 X2 X0)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((\neg v1_rat_1 X0) \Leftrightarrow (X0 \in k1_borsuk_5)) \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (k2_rcomp_1 X0 X1 = k4_xxreal_1 X0 X1) \tag{4}$$

Assume the following.

$$m1_subset_1 k1_borsuk_5 (k1_zfmisc_1 k1_numbers) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \tag{6}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (k3_borsuk_5 X0 X1 = k9_subset_1 k1_numbers k1_borsuk_5 (k2_rcomp_1 X0 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k9_subset_1 X0 X2 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1_rat_1 X0) \Rightarrow (v1_xreal_0 X0) \quad (11)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow ((X2 \in k3_borsuk_5 X0 X1) \Leftrightarrow ((\neg v1_rat_1 X2) \wedge ((\neg r1_xxreal_0 X2 X0) \wedge (\neg r1_xxreal_0 X1 X2)))))$$