

t30_csspace4

(TMXn8DvxYpybgBn7dZ8LWVDozW1KCowNgVd)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k2_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_comseq_2 : \iota \Rightarrow o$ be given. Let $k55_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers)))))) \Rightarrow \\ & ((v1_comseq_2 (k55_valued_1 k5_numbers k2_numbers X0)) \Rightarrow (v1_comseq_2 \\ & X0)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers)))))) \Rightarrow \\ & ((v1_comseq_2 X0) \Rightarrow (v1_comseq_2 (k55_valued_1 k5_numbers k2_numbers \\ & X0))) \end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers)))))) \Rightarrow \\ & ((v1_comseq_2 X0) \Leftrightarrow (v1_comseq_2 (k55_valued_1 k5_numbers k2_numbers \\ & X0))) \end{aligned}$$