

## t30\_domain\_1

(TMQY3u1upEtuYUvSBdfRDMXy9tHjVYxkPtc)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. \neg(X0 \in X1) \wedge (v1\_xboole\_0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg(X0 \in X1) \wedge ((m1\_subset\_1 X1 (k1\_zfmisc\_1 X2)) \wedge (v1\_xboole\_0 X2)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X2))) \Rightarrow (m1\_subset\_1 X0 X2) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1\_subset\_1 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (k7\_subset\_1 X0 X1 X2 = k4\_xboole\_0 X1 X2) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k4\_xboole\_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \quad (7)$$

### Theorem 1

$$\begin{aligned} & \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow (k7\_subset\_1 X0 X1 X2 = \text{ReplSep} (\text{toset} (\lambda X3 : \iota. m1\_subset\_1 X3 X0)) (\lambda X3 : \iota. (X3 \in X1) \wedge (\neg X3 \in X2)) (\lambda X3 : \iota. X3)))) \end{aligned}$$