

t30_euclid

(TMP5zX34YDRZG5x7Y92ZJsM4cx7YvS8d2cr)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k1_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k11_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rltopsp1 : \iota \Rightarrow o$ be given. Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (k11_binop_2 X0 X1 = k3_xcmplx_0 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (v7_ordinal1 X0) \Rightarrow & ((v2_pre_topc (k15_euclid X0)) \wedge \\ & ((v13_algstr_0 (k15_euclid X0)) \wedge ((v2_rlvect_1 (k15_euclid X0)) \wedge \\ & ((v3_rlvect_1 (k15_euclid X0)) \wedge ((v4_rlvect_1 (k15_euclid X0)) \wedge \\ & ((v5_rlvect_1 (k15_euclid X0)) \wedge ((v6_rlvect_1 (k15_euclid X0)) \wedge \\ & ((v7_rlvect_1 (k15_euclid X0)) \wedge ((v8_rlvect_1 (k15_euclid X0)) \wedge \\ & (v5_rltopsp1 (k15_euclid X0)))))))))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow ((-v2_struct_0 (k15_euclid X0)) \wedge (v5_rltopsp1 (k15_euclid X0))) \quad (3)$$

Assume the following.

$$\forall X0. (l1_rltopsp1 X0) \Rightarrow ((l1_rlvect_1 X0) \wedge (l1_pre_topc X0)) \quad (4)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow ((v5_rltopsp1 (k15_euclid X0)) \wedge (l1_rltopsp1 (k15_euclid X0))) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_rlvect_1 X0)) \Rightarrow ((v7_rlvect_1 \\ & X0) \Leftrightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow \\ & (\forall X3.(m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow (k1_rlvect_1 X0 \\ & X3 (k3_xcmplx_0 X1 X2) = k1_rlvect_1 X0 (k1_rlvect_1 X0 X3 X2) X1)))))) \end{aligned} \tag{6}$$

Theorem 1

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 \\ & (k15_euclid X0))) \Rightarrow (\forall X2.(v1_xreal_0 X2) \Rightarrow (\forall X3.(\\ & v1_xreal_0 X3) \Rightarrow (k1_rlvect_1 (k15_euclid X0) X1 (k11_binop_2 X2 \\ & X3) = k1_rlvect_1 (k15_euclid X0) (k1_rlvect_1 (k15_euclid X0) \\ & X1 X3) X2)))))) \end{aligned}$$