

## t30\_finseq\_3

(TMM8qghH2FobZVREBAinxHAJf26K5dFHzbW)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((r1\_xxreal\_0 X0 X1) \Leftrightarrow (r1\_tarski (k2\_finseq\_1 X0) (k2\_finseq\_1 X1)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \quad (2)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (4)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1) \wedge (v3\_ordinal1 k4\_ordinal1) \quad (5)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (6)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (m2\_subset\_1 (k3\_finseq\_1 X0) k1\_numbers k5\_numbers) \quad (7)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (\forall X1.(m2\_subset\_1 X1 k1\_numbers k5\_numbers) \Rightarrow ((X1 = k3\_finseq\_1 X0) \Leftrightarrow (k2\_finseq\_1 X1 = k9\_xtuple\_0 X0))) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (9)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow ((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0)))) \quad (10)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (v1\_xboole\_0 X1)) \quad (11)$$

**Theorem 1**

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 X1))) \Rightarrow ((r1\_xxreal\_0 (k3\_finseq\_1 X0) (k3\_finseq\_1 X1)) \Leftrightarrow (r1\_tarski (k1\_relset\_1 k5\_numbers X0) (k1\_relset\_1 k5\_numbers X1))))$$