

t30\_funct\_2  
(TMSghcWwTobTdjRcnjergfkyPdVMjxLKu65)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_partfun1 : \iota \Rightarrow \iota$  be given. Let  $v2\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k2\_funct\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k3\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_relat\_1 : \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.(( \\ v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (((v2\_funct\_1 X0) \wedge ((k10\_xtuple\_0 \\ X0 = k9\_xtuple\_0 X1) \wedge (k3\_relat\_1 X0 X1 = k4\_relat\_1 (k9\_xtuple\_0 \\ X0)))) \Rightarrow (X1 = k2\_funct\_1 X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. \forall X3. ((m1\_subset\_1 X2 \\ (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ (k2\_zfmisc\_1 X0 X1)))) \Rightarrow ((r2\_relset\_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0. k6\_partfun1 X0 = k4\_relat\_1 X0 \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v5\_relat\_1 X1 X0)) \Rightarrow ( \\ k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \tag{4}$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow ( \\ k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \tag{5}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.\forall X4.\forall X5. \\ & (((v1\_funct\_1 X4)\wedge(m1\_subset\_1 X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 X1))))\wedge((v1\_funct\_1 X5)\wedge(m1\_subset\_1 X5 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X2 X3))))\Rightarrow(k1\_partfun1 X0 X1 X2 X3 X4 X5 = k3\_relat\_1 X4 X5) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1\_partfun1 (k6\_partfun1 X0) X0)\wedge(m1\_subset\_1 (k6\_partfun1 X0) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X0))) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.\forall X4.\forall X5. \\ & (((v1\_funct\_1 X4)\wedge(m1\_subset\_1 X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 X1))))\wedge((v1\_funct\_1 X5)\wedge(m1\_subset\_1 X5 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X2 X3))))\Rightarrow((v1\_funct\_1 (k1\_partfun1 X0 X1 X2 X3 X4 X5))\wedge(m1\_subset\_1 \\ & (k1\_partfun1 X0 X1 X2 X3 X4 X5) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X3)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1)))\Rightarrow(((X1\neq k1\_xboole\_0)\Rightarrow((v1\_funct\_2 X2 X0 \\ & X1)\Leftrightarrow(X0 = k1\_relset\_1 X0 X2)))\wedge((X1 = k1\_xboole\_0)\Rightarrow((v1\_funct\_2 \\ & X2 X0 X1)\Leftrightarrow(X2 = k1\_xboole\_0)))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (11)$$

### Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1\_funct\_1 X2)\wedge((v1\_funct\_2 \\ & X2 X0 X1)\wedge(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))\Rightarrow \\ & (\forall X3.((v1\_funct\_1 X3)\wedge((v1\_funct\_2 X3 X1 X0)\wedge(m1\_subset\_1 \\ & X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 X0))))\Rightarrow(((k2\_relset\_1 X1 X2 = \\ & X1)\wedge((r2\_relset\_1 X0 X0 (k1\_partfun1 X0 X1 X1 X0 X2 X3) (k6\_partfun1 \\ & X0))\wedge(v2\_funct\_1 X2)))\Rightarrow((X0 = k1\_xboole\_0)\vee((X1 = k1\_xboole\_0)\vee \\ & (X3 = k2\_funct\_1 X2)))) \end{aligned}$$