

t30_group_2
(TMK2VRswZDVCyYoQZdNjaR5RapgvUeiz38j)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l3_algstr_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v3_group_1 : \iota \Rightarrow o$ be given. Let $k2_group_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_group_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_group_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l3_algstr_0 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X3. \\ & (m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((v3_group_1 \\ & X0) \Rightarrow (k2_group_2 X0 (k2_group_2 X0 X1 X2) X3 = k2_group_2 X0 X1 (k2_group_2 \\ & X0 X2 X3)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (u1_struct_0 X0)) \tag{2}$$

Assume the following.

$$\forall X0.(l3_algstr_0 X0) \Rightarrow (l1_struct_0 X0) \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (m1_subset_1 (k6_domain_1 X0 X1) (k1_zfmisc_1 X0)) \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l3_algstr_0 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 \\ & (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k5_group_2 X0 X1 X2 = k2_group_2 \\ & X0 X2 (k6_domain_1 (u1_struct_0 X0) X1)))) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge (l3_algstr_0 X0)) \Rightarrow (\forall X1. \\
& (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 \\
& (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (k4_group_2 X0 X1 X2 = k2_group_2 \\
& X0 (k6_domain_1 (u1_struct_0 X0) X1) X2)))
\end{aligned} \tag{6}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge (l3_algstr_0 X0)) \Rightarrow (\forall X1. \\
& (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2. \\
& (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X3. \\
& (m1_subset_1 X3 (u1_struct_0 X0)) \Rightarrow ((v3_group_1 X0) \Rightarrow (k2_group_2 \\
& X0 (k5_group_2 X0 X3 X1) X2 = k2_group_2 X0 X1 (k4_group_2 X0 X3 X2))))))
\end{aligned}$$