

t30_helly

(TMS7bVa5AsybrgwgbGF9mpXCJuKdCJrm2Zv)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $v3_glib_002 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_glib_000 : \iota \Rightarrow \iota$ be given. Let $k13_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_helly : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_glib_001 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m3_glib_001 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_glib_001 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_glib_000 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v5_glib_001 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ & X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 (k6_glib_000 X0)) \Rightarrow (k13_glib_001 X0 (k1_glib_001 X0 X1) = k6_domain_1 \\ & (k6_glib_000 X0) X1)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ & X0) \wedge ((v1_finset_1 X0) \wedge ((v1_glib_000 X0) \wedge (v3_glib_002 X0)))))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (k6_glib_000 X0)) \Rightarrow (\forall X2.(m1_subset_1 \\ & X2 (k6_glib_000 X0)) \Rightarrow ((k3_glib_001 X0 (k2_helly X0 X1 X2) = X1) \wedge \\ & (k4_glib_001 X0 (k2_helly X0 X1 X2) = X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 (k6_glib_000 X0)) \Rightarrow ((k3_finseq_1 (k1_glib_001 X0 X1) = np_1) \wedge \\ ((k1_funct_1 (k1_glib_001 X0 X1) np_1 = X1) \wedge ((k3_glib_001 X0 (\\ k1_glib_001 X0 X1) = X1) \wedge ((k4_glib_001 X0 (k1_glib_001 X0 X1) = X1) \wedge \\ (r1_glib_001 X0 X1 X1 (k1_glib_001 X0 X1))))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.(m3_glib_001 \\ X1 X0) \Rightarrow ((v3_glib_001 X1 X0) \Leftrightarrow (\exists X2.(m1_subset_1 X2 (k6_glib_000 \\ X0)) \wedge (r2_relset_1 k5_numbers (k2_xboole_0 (k6_glib_000 X0) (\\ k7_glib_000 X0)) X1 (k1_glib_001 X0 X2)))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. \forall X3. ((m1_subset_1 X2 \\ (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \wedge (m1_subset_1 X3 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1)))) \Rightarrow ((r2_relset_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow \\ (k6_domain_1 X0 X1 = k1_tarski X1) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge \\ ((v1_funct_1 X0) \wedge ((v1_finset_1 X0) \wedge ((v1_glib_000 X0) \wedge (v3_glib_002 \\ X0)))))) \wedge (m1_subset_1 X1 (k6_glib_000 X0))) \Rightarrow ((v3_glib_001 (\\ k2_helly X0 X1 X1) X0) \wedge (v5_glib_001 (k2_helly X0 X1 X1) X0)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\neg v1_xboole_0 (\\ k6_glib_000 X0)) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge ((v1_finset_1 X0) \wedge (v1_glib_000 X0)))))) \Rightarrow (\forall X1.(m3_glib_001 \\ X1 X0) \Rightarrow (m2_finseq_1 X1 (k2_xboole_0 (k6_glib_000 X0) (k7_glib_000 \\ X0)))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge((v1_glib_000 X0)\wedge(v3_glib_002 X0))))))\wedge((m1_subset_1 X1 (k6_glib_000 X0))\wedge(m1_subset_1 X2 (k6_glib_000 X0))))\Rightarrow((v5_glib_001 (k2_helly X0 X1 X2) X0)\wedge(m3_glib_001 (k2_helly X0 X1 X2) X0)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge(v1_glib_000 X0)))))\wedge(m1_subset_1 X1 (k6_glib_000 X0)))\Rightarrow(m3_glib_001 (k1_glib_001 X0 X1) X0) \quad (12)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge((v1_glib_000 X0)\wedge(v3_glib_002 X0))))))\Rightarrow(\forall X1.(m1_subset_1 X1 (k6_glib_000 X0))\Rightarrow(k13_glib_001 X0 (k2_helly X0 X1 X1) = k1_tarski X1))$$