

t30_nat_d (TMWqCxxajaijMsDcPLDB-
JqB92DMA2Tu6jP3)

October 27, 2020

Let $v1_int_1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k3_int_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_int_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k6_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_nat_d : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k1_int_2 : \iota \Rightarrow \iota$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 X1) \wedge ((\neg v3_xxreal_0 X1) \wedge (\neg v2_xxreal_0 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 X0 k6_numbers) \Rightarrow (k6_nat_d X1 X0 = k6_nat_d X0 (k4_nat_d X1 X0)))) \quad (4)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (5)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(k6_nat_d X0 X1 = k3_int_2 X0 X1) \quad (7)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(k4_nat_d X0 X1 = k6_int_1 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.(v1_int_1 X0)\Rightarrow(k1_int_2 X0 = k16_complex1 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0)\Rightarrow(k4_xcmplx_0 (k4_xcmplx_0 X0) = X0) \quad (11)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (12)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow((v1_xcmplx_0 (k4_xcmplx_0 X0))\wedge(v1_xreal_0 (k4_xcmplx_0 X0))) \quad (13)$$

Assume the following.

$$\forall X0.((\neg v3_xreal_0 X0)\wedge(v1_xreal_0 X0))\Rightarrow((v1_xcmplx_0 (k4_xcmplx_0 X0))\wedge(\neg v2_xreal_0 (k4_xcmplx_0 X0))) \quad (14)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(m1_subset_1 (k4_nat_d X0 X1) k5_numbers) \quad (16)$$

Assume the following.

$$\forall X0.(v1_int_1 X0)\Rightarrow(m1_subset_1 (k1_int_2 X0) k5_numbers) \quad (17)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \Rightarrow (k16_complex1 X0 = X0)) \wedge ((\neg r1_xxreal_0 k6_numbers X0) \Rightarrow (k16_complex1 X0 = k4_xcmplx_0 X0))) \quad (18)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (19)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (20)$$

Assume the following.

$$\forall X0.(v1_int_1 X0) \Rightarrow (v1_xreal_0 X0) \quad (21)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k5_numbers) \Rightarrow (\neg v3_xxreal_0 X0) \quad (22)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (23)$$

Theorem 1

$$\forall X0.(v1_int_1 X0) \Rightarrow (\forall X1.(v1_int_1 X1) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \Rightarrow ((r1_xxreal_0 X1 k6_numbers) \vee (k3_int_2 X0 X1 = k3_int_2 X1 (k6_int_1 X0 X1))))))$$