

t30_procal_1
(TMb9L1i2o3QjLMMfiYSWFSR3jUSMCSFEQuQ)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $k10_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_cqc_the1 : \iota \Rightarrow \iota$ be given. Let $k8_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (k8_cqc_lang\ X0\ (k9_cqc_lang\ X0\ X1\ X2)\ (k9_cqc_lang \\ X0\ X2\ X1) \in k4_cqc_the1\ X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (k8_cqc_lang\ X0\ (k8_cqc_lang\ X0\ X1\ X2)\ (k8_cqc_lang \\ X0\ (k8_cqc_lang\ X0\ X2\ X1)\ (k10_cqc_lang\ X0\ X1\ X2)) \in k4_cqc_the1\ X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X3.(m2_subset_1\ X3\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (((k8_cqc_lang\ X0\ X1\ (k8_cqc_lang\ X0\ X2\ X3) \in \\ k4_cqc_the1\ X0) \wedge ((X2 \in k4_cqc_the1\ X0) \wedge (X1 \in k4_cqc_the1\ X0))) \Rightarrow \\ (X3 \in k4_cqc_the1\ X0)))))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)) \Rightarrow (m2_subset_1\ (k4_qc_lang2\ X0\ X1\ X2)\ (k9_qc_lang1 \\ X0)\ (k3_cqc_lang\ X0)))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Leftrightarrow(m1_subset_1 X2 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0)\wedge((m1_subset_1 X1 (k3_cqc_lang X0))\wedge(m1_subset_1 X2 (k3_cqc_lang X0))))\Rightarrow(k10_cqc_lang X0 X1 X2 = k4_qc_lang2 X0 X1 X2) \quad (6)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(\neg v1_xboole_0 (k3_cqc_lang X0)) \quad (7)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(\neg v1_xboole_0 (k9_qc_lang1 X0)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0)\wedge((m1_subset_1 X1 (k3_cqc_lang X0))\wedge(m1_subset_1 X2 (k3_cqc_lang X0))))\Rightarrow(m2_subset_1 (k9_cqc_lang X0 X1 X2) (k9_qc_lang1 X0) (k3_cqc_lang X0)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_qc_lang1 X0)\wedge((m1_subset_1 X1 (k3_cqc_lang X0))\wedge(m1_subset_1 X2 (k3_cqc_lang X0))))\Rightarrow(m2_subset_1 (k8_cqc_lang X0 X1 X2) (k9_qc_lang1 X0) (k3_cqc_lang X0)) \quad (10)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(m1_subset_1 (k3_cqc_lang X0) (k1_zfmisc_1 (k9_qc_lang1 X0))) \quad (11)$$

Theorem 1

$$\forall X0.(m1_qc_lang1 X0)\Rightarrow(\forall X1.(m2_subset_1 X1 (k9_qc_lang1 X0) (k3_cqc_lang X0))\Rightarrow(\forall X2.(m2_subset_1 X2 (k9_qc_lang1 X0) (k3_cqc_lang X0))\Rightarrow(k10_cqc_lang X0 (k9_cqc_lang X0 X1 X2) (k9_cqc_lang X0 X2 X1) \in k4_cqc_the1 X0)))$$