

t30_rusub_3

(TMRe3RUZhbAN8QYfMMBQvLvN1GAef77ye7p)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v2_bhsp_1 : \iota \Rightarrow o$ be given. Let $l1_bhsp_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_struct_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rusub_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k1_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\ & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge ((v2_bhsp_1 \\ & X0) \wedge (l1_bhsp_1 X0)))))))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 \\ & X0)) \Rightarrow (\forall X2.(r1_struct_0 (k1_rusub_3 X0 (k6_domain_1 (u1_struct_0 \\ & X0) X1)) X2) \Leftrightarrow (\exists X3.(m1_subset_1 X3 k1_numbers) \wedge (X2 = k1_rlvect_1 \\ & X0 X1 X3)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.(l1_bhsp_1 X0) \Rightarrow (l1_rlvect_1 X0) \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_rlvect_1 X0)) \Rightarrow ((v8_rlvect_1 \\ & X0) \Leftrightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (k1_rlvect_1 \\ & X0 X1 np_1 = X1))) \end{aligned} \tag{4}$$

Theorem 1

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (v13_algstr_0 X0) \wedge (v2_rlvect_1 \\ & X0) \wedge (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge (v7_rlvect_1 X0) \wedge (v8_rlvect_1 X0) \wedge (v2_bhsp_1 \\ & X0) \wedge (l1_bhsp_1 X0)))))) \Rightarrow (\forall X1. (m1_subset_1 X1 (u1_struct_0 \\ & X0)) \Rightarrow (r1_struct_0 (k1_rusub_3 X0 (k6_domain_1 (u1_struct_0 X0) \\ & X1)) X1)) \end{aligned}$$