

t30_rvsum_2 (TMNXeXp-
BRZ1umXD6a889otTnpzTtw9VL5zR)

October 27, 2020

Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k16_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_finsop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_numbers : \iota$ be given. Let $k17_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $k27_binop_2 : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow \\ & (\forall X2. ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 (k2_zfmisc_1 X0 \\ & X0) X0) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 \\ & X0 X0) X0)))))) \Rightarrow (k1_finsop_1 X0 (k12_finseq_1 X0 X1) X2 = X1))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0. k9_finseq_1 X0 = k5_finseq_1 X0 \quad (4)$$

Assume the following.

$$\forall X0. (m1_finseq_1 X0 k2_numbers) \Rightarrow (k17_rvsum_1 X0 = k16_rvsum_1 X0) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 \ k2_numbers \quad (7)$$

Assume the following.

$$\begin{aligned} & (v1_funct_1 \ k27_binop_2) \wedge ((v1_funct_2 \ k27_binop_2 \ (k2_zfmisc_1 \\ & k2_numbers \ k2_numbers) \ k2_numbers) \wedge (m1_subset_1 \ k27_binop_2 \\ & (k1_zfmisc_1 \ (k2_zfmisc_1 \ (k2_zfmisc_1 \ k2_numbers \ k2_numbers) \\ & k2_numbers)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 \ X0) \wedge (m1_subset_1 \ X1 \ X0)) \Rightarrow \\ & (m2_finseq_1 \ (k12_finseq_1 \ X0 \ X1) \ X0) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0. (v1_xcmplx_0 \ X0) \Leftrightarrow (X0 \in k2_numbers) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0. (m2_finseq_1 \ X0 \ k2_numbers) \Rightarrow (k17_rvsum_1 \ X0 = k1_finsop_1 \\ & k2_numbers \ X0 \ k27_binop_2) \end{aligned} \quad (11)$$

Theorem 1 $\forall X0. (v1_xcmplx_0 \ X0) \Rightarrow (k16_rvsum_1 \ (k9_finseq_1 \ X0) = X0).$