

t30_scmfsa_2

(TMWAh73pxptnemVHHDg4sh75oLjqEwTG7J8)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_compos_1 : \iota \Rightarrow \iota$ be given. Let $k1_scmfsa_2 : \iota$ be given. Let $k2_compos_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_ami_2 : \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k6_scmfsa_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_ami_3 : \iota$ be given. Let $k2_ami_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_8 : \iota$ be given. Let $k2_scmfsa_2 : \iota$ be given. Let $k2_scm_inst : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_compos_0 : \iota \Rightarrow o$ be given. Let $k4_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_ami_2 : \iota$ be given. Let $l1_compos_1 : \iota \Rightarrow o$ be given. Let $v2_compos_0 : \iota \Rightarrow o$ be given. Let $v3_compos_0 : \iota \Rightarrow o$ be given. Let $v5_compos_0 : \iota \Rightarrow o$ be given. Let $l1_extpro_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $l1_memstr_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_extpro_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_3 : \iota$ be given. Let $np_2 : \iota$ be given. Let $k1_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_compos_1 k1_ami_3)) \Rightarrow & (\neg(k2_compos_0 \\ & (u1_compos_1 k1_ami_3) X0 = np_1) \wedge (\forall X1.((v1_ami_2 X1) \wedge \\ & (m1_subset_1 X1 (u1_struct_0 k1_ami_3))) \Rightarrow (\forall X2.((v1_ami_2 \\ & X2) \wedge (m1_subset_1 X2 (u1_struct_0 k1_ami_3))) \Rightarrow (X0 \neq k2_ami_3 X1 \\ & X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_compos_1 k1_scmfsa_2)) \Rightarrow & ((r1_xxreal_0 \\ & (k2_compos_0 (u1_compos_1 k1_scmfsa_2) X0) np_8) \Rightarrow (m1_subset_1 \\ & X0 (u1_compos_1 k1_ami_3))) \end{aligned} \quad (3)$$

Assume the following.

$$r1_xxreal_0 np_1 np_8 \quad (4)$$

Assume the following.

$$k2_scmfsa_2 = k2_scm_inst \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (((\neg v1_xboole_0 X0) \wedge (v1_compos_0 X0)) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k2_compos_0 X0 X1 = k4_xtuple_0 X1) \quad (6)$$

Assume the following.

$$k2_ami_2 = k2_scm_inst \quad (7)$$

Assume the following.

$$\forall X0. (l1_compos_1 X0) \Rightarrow ((v1_compos_0 (u1_compos_1 X0)) \wedge ((v2_compos_0 (u1_compos_1 X0)) \wedge (v3_compos_0 (u1_compos_1 X0)) \wedge (v5_compos_0 (u1_compos_1 X0)))) \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. (l1_extpro_1 X1 X0) \Rightarrow ((l1_memstr_0 X1 X0) \wedge (l1_compos_1 X1)) \quad (9)$$

Assume the following.

$$m1_subset_1 k2_scmfsa_2 (k1_zfmisc_1 (u1_struct_0 k1_scmfsa_2)) \quad (10)$$

Assume the following.

$$(v1_extpro_1 k1_scmfsa_2 np_3) \wedge (l1_extpro_1 k1_scmfsa_2 np_3) \quad (11)$$

Assume the following.

$$(v1_extpro_1 k1_ami_3 np_2) \wedge (l1_extpro_1 k1_ami_3 np_2) \quad (12)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_ami_2 X0) \wedge (m1_subset_1 X0 (u1_struct_0 k1_scmfsa_2))) \Rightarrow \\ & (\forall X1. ((v1_ami_2 X1) \wedge (m1_subset_1 X1 (u1_struct_0 k1_scmfsa_2))) \Rightarrow \\ & (\forall X2. (m1_subset_1 X2 (u1_compos_1 k1_scmfsa_2)) \Rightarrow ((X2 = \\ & k6_scmfsa_2 X0 X1) \Leftrightarrow (\exists X3. ((v1_ami_2 X3) \wedge (m1_subset_1 X3 \\ & (u1_struct_0 k1_ami_3)))) \wedge (\exists X4. ((v1_ami_2 X4) \wedge (m1_subset_1 \\ & X4 (u1_struct_0 k1_ami_3)))) \wedge ((X0 = X3) \wedge ((X1 = X4) \wedge (X2 = k2_ami_3 \\ & X3 X4)))))) \quad (13) \end{aligned}$$

Assume the following.

$$\forall X0. k4_xtuple_0 X0 = k1_xtuple_0 (k1_xtuple_0 X0) \quad (14)$$

Assume the following.

$$\forall X0. (v1_ami_2 X0) \Leftrightarrow (X0 \in k2_ami_2) \quad (15)$$

Assume the following.

$$\forall X0. (v5_compos_0 X0) \Rightarrow (\neg v1_xboole_0 X0) \quad (16)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (u1_compos_1 k1_scmfsa_2)) \Rightarrow & (\neg(k2_compos_0 \\ & (u1_compos_1 k1_scmfsa_2) X0 = np_1) \wedge (\forall X1.((v1_ami_2 \\ & X1) \wedge (m1_subset_1 X1 (u1_struct_0 k1_scmfsa_2))) \Rightarrow (\forall X2. \\ & ((v1_ami_2 X2) \wedge (m1_subset_1 X2 (u1_struct_0 k1_scmfsa_2))) \Rightarrow \\ & (X0 \neq k6_scmfsa_2 X1 X2)))) \end{aligned}$$