

t30_xreal_1

(TMY7EyU82UxmN1w45sFTzy2v2kWwAR7aR3p)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (X0 = k6_xcmplx_0 (k2_xcmplx_0 X0 X1) X1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (r1_xxreal_0 k6_numbers (k6_xcmplx_0 X1 X0)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k2_xcmplx_0 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 X1 (k2_xcmplx_0 X0 X1))))$$