

t30_yellow18

(TMHB2VXFd4g9NdSJcPk2PCcK4PGABoS2py7)

October 27, 2020

Let $k1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 \\ X2)) \Rightarrow ((X2 \in k1_funct_2 X0 X1) \Rightarrow ((k9_xtuple_0 X2 = X0) \wedge (r1_tarski \\ (k10_xtuple_0 X2) X1))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (X2 = k1_funct_2 X0 X1) \Leftrightarrow (\forall X3. \\ (X3 \in X2) \Leftrightarrow (\exists X4. ((v1_relat_1 X4) \wedge (v1_funct_1 X4)) \wedge ((X3 = \\ X4) \wedge ((k9_xtuple_0 X4 = X0) \wedge (r1_tarski (k10_xtuple_0 X4) X1)))))) \end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (X2 \in k1_funct_2 X0 X1) \Leftrightarrow (((v1_relat_1 \\ X2) \wedge (v1_funct_1 X2)) \wedge ((k9_xtuple_0 X2 = X0) \wedge (r1_tarski (k10_xtuple_0 \\ X2) X1))) \end{aligned}$$