

t31_bhsp_3

(TMFaye9fpBkM1vECszvwreWCSvLFU3oYmpi)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v2_bhsp_1 : \iota \Rightarrow o$ be given. Let $l1_bhsp_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_valued_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k9_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $m2_valued_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $l2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_algstr_0 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. ((v1_xcmplx_0 X0) \wedge ((v1_xcmplx_0 \\ X1) \wedge (v1_xcmplx_0 X2))) \Rightarrow (k2_xcmplx_0 (k2_xcmplx_0 X0 X1) X2 = k2_xcmplx_0 \\ X0 (k2_xcmplx_0 X1 X2)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{2}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0))))))\wedge(v7_ordinal1 X2))\Rightarrow(k1_valued_0 X0 X1 X2 = k9_nat_1 X1 X2) \quad (5)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (6)$$

Assume the following.

$$v3_membered k1_numbers \quad (7)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0)\wedge(l1_struct_0 X0))\Rightarrow(\neg v1_xboole_0 (u1_struct_0 X0)) \quad (8)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v7_ordinal1 (k2_xcmplx_0 X0 X1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers X0)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0))))))\Rightarrow(\forall X2.(m2_valued_0 X2 X0 X1)\Rightarrow((v1_funct_1 X2)\wedge((v1_funct_2 X2 k5_numbers X0)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (11)$$

Assume the following.

$$\forall X0.(l2_algstr_0 X0)\Rightarrow((l2_struct_0 X0)\wedge(l1_algstr_0 X0)) \quad (12)$$

Assume the following.

$$\forall X0.(l1_rlvect_1 X0)\Rightarrow(l2_algstr_0 X0) \quad (13)$$

Assume the following.

$$\forall X0.(l1_bhsp_1 X0)\Rightarrow(l1_rlvect_1 X0) \quad (14)$$

Assume the following.

$$\forall X0.(l1_algstr_0 X0) \Rightarrow (l1_struct_0 X0) \quad (15)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.(((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge \\ ((v1_funct_1 X0) \wedge (v1_partfun1 X0 k5_numbers)))) \wedge (v7_ordinal1 \\ X1)) \Rightarrow ((v1_relat_1 (k9_nat_1 X0 X1)) \wedge ((v4_relat_1 (k9_nat_1 X0 \\ X1) k5_numbers) \wedge ((v1_funct_1 (k9_nat_1 X0 X1)) \wedge (v1_partfun1 \\ (k9_nat_1 X0 X1) k5_numbers)))))) \end{aligned} \quad (16)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (17)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0) \wedge (((v1_funct_1 \\ X1) \wedge ((v1_funct_2 X1 k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ (k2_zfmisc_1 k5_numbers X0)))))) \wedge (v7_ordinal1 X2))) \Rightarrow (m2_valued_0 \\ (k1_valued_0 X0 X1 X2) X0 X1) \end{aligned} \quad (18)$$

Assume the following.

$$\begin{aligned} \forall X0.(((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ X0) \wedge (v1_partfun1 X0 k5_numbers)))) \Rightarrow (\forall X1.(v7_ordinal1 \\ X1) \Rightarrow (\forall X2.(((v1_relat_1 X2) \wedge ((v4_relat_1 X2 k5_numbers) \wedge \\ ((v1_funct_1 X2) \wedge (v1_partfun1 X2 k5_numbers)))))) \Rightarrow ((X2 = k9_nat_1 \\ X0 X1) \Leftrightarrow (\forall X3.(v7_ordinal1 X3) \Rightarrow (k1_funct_1 X2 X3 = k1_funct_1 \\ X0 (k2_xcmplx_0 X3 X1)))))) \end{aligned} \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (20)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (21)$$

Assume the following.

$$\forall X0.(v3_membered X0) \Rightarrow (v1_membered X0) \quad (22)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (23)$$

Assume the following.

$$\forall X0.\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v1_funct_2 X2 X0 X1)\Rightarrow(v1_partfun1 X2 X0))) \quad (24)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xreal_0 X0) \quad (25)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (26)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (27)$$

Assume the following.

$$\forall X0.(v1_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_membered X1)) \quad (28)$$

Assume the following.

$$\forall X0.(v1_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v1_xcmplx_0 X1)) \quad (29)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0)\wedge((v13_algstr_0 X0)\wedge((v2_rlvect_1 X0)\wedge((v3_rlvect_1 X0)\wedge((v4_rlvect_1 X0)\wedge((v5_rlvect_1 X0)\wedge \\ & ((v6_rlvect_1 X0)\wedge((v7_rlvect_1 X0)\wedge((v8_rlvect_1 X0)\wedge((v2_bhsp_1 X0)\wedge(l1_bhsp_1 X0))))))))))\Rightarrow(\forall X1.((v1_funct_1 X1)\wedge \\ & ((v1_funct_2 X1 k5_numbers (u1_struct_0 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (u1_struct_0 X0))))))\Rightarrow \\ & (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers)\Rightarrow(\forall X3. \\ & (m2_subset_1 X3 k1_numbers k5_numbers)\Rightarrow(k1_valued_0 (u1_struct_0 X0) (k1_valued_0 (u1_struct_0 X0) X1 X2) X3 = k1_valued_0 (u1_struct_0 X0) X1 (k2_nat_1 X2 X3)))))) \end{aligned}$$