

t31_enumset1 (TMUddRdnSh- froWqVgTZpCzEdE1Rv2gaQ9DT)

October 27, 2020

Let $k2_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. k1_enumset1 X0 X1 X2 = k2_xboole_0 (k1_tarski X0) (k2_tarski X1 X2) \quad (1)$$

Assume the following.

$$\forall X0. k2_tarski X0 X0 = k1_tarski X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. k2_enumset1 X0 X1 X2 X3 = k2_xboole_0 (k2_tarski X0 X1) (k2_tarski X2 X3) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. \forall X2. k2_enumset1 X0 X0 X1 X2 = k1_enumset1 X0 X1 X2$$