

t31_finseq_2

(TMX4VXX8rPbBqRVF7r3KjYJSEbNCfgu14tX)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow (\\ & (X1 = k2_finseq_1 X0) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge ((v1_funct_1 \\ & X2) \wedge (v1_finseq_1 X2)))) \Rightarrow (\forall X3.(m2_finseq_1 X3 X1) \Rightarrow ((r1_xxreal_0 \\ & X0 (k3_finseq_1 X2)) \Rightarrow ((v1_relat_1 (k3_relat_1 X3 X2)) \wedge ((v1_funct_1 \\ & (k3_relat_1 X3 X2)) \wedge (v1_finseq_1 (k3_relat_1 X3 X2)))))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.(v1_relat_1 X1) \Rightarrow (r1_tarski (k10_xtuple_0 (k3_relat_1 X0 X1)) (k10_xtuple_0 X1))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((r1_tarski X0 X1) \wedge (r1_tarski X1 X2)) \Rightarrow (r1_tarski X0 X2) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.\forall X4.\forall X5. \\ & (((v1_funct_1 X4) \wedge (m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X0 X1)))) \wedge ((v1_funct_1 X5) \wedge (m1_subset_1 X5 (k1_zfmisc_1 (k2_zfmisc_1 \\ & X2 X3)))))) \Rightarrow (k1_partfun1 X0 X1 X2 X3 X4 X5 = k3_relat_1 X4 X5) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k3_relat_1 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1)))\Rightarrow((m1_finseq_1 X1 X0)\Leftrightarrow(r1_tarski (k10_xtuple_0 X1) X0)) \quad (9)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.(\neg v1_xboole_0 X2)\Rightarrow((X1 = k2_finseq_1 X0)\Rightarrow(\forall X3.(m2_finseq_1 X3 X2)\Rightarrow(\forall X4.(m2_finseq_1 X4 X1)\Rightarrow((r1_xxreal_0 X0 (k3_finseq_1 X3))\Rightarrow(m2_finseq_1 (k1_partfun1 k5_numbers X1 k5_numbers X2 X4 X3) X2))))))))))$$