

t31_finseq_3 (TMHUUVfD- jpBmNN9Hw9o7DroFNAVNWkijHNn)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1\ X0) \Rightarrow (\forall X1.(v7_ordinal1\ X1) \Rightarrow ((r1_xxreal_0\ X0\ X1) \Leftrightarrow (r1_tarski\ (k2_finseq_1\ X0)\ (k2_finseq_1\ X1)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1\ X0\ (k1_zfmisc_1\ X1)) \Leftrightarrow (r1_tarski\ X0\ X1) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0) \Rightarrow (\forall X1.(v7_ordinal1\ X1) \Rightarrow ((X0 \in k2_finseq_1\ X1) \Leftrightarrow ((r1_xxreal_0\ np_1\ X0) \wedge (r1_xxreal_0\ X0\ X1)))) \quad (3)$$

Assume the following.

$$((v2_xxreal_0\ np_1) \wedge (m2_subset_1\ np_1\ k1_numbers\ k5_numbers)) \wedge ((m1_subset_1\ np_1\ k5_numbers) \wedge (m1_subset_1\ np_1\ k1_numbers)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.r1_tarski\ X0\ X0 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (7)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (8)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m1_subset_1 (k4_finseq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (10)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. (X1 = k10_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. (X3 \in k9_xtuple_0 X0) \wedge (X2 = k1_funct_1 X0 X3)))) \quad (11)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow ((v1_finseq_1 X0) \Leftrightarrow (\exists X1. (v7_ordinal1 X1) \wedge (k9_xtuple_0 X0 = k2_finseq_1 X1))) \quad (12)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Leftrightarrow (X0 \in k4_ordinal1) \quad (13)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (14)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow ((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0)))) \quad (15)$$

Theorem 1

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (\forall X1. (X1 \in k10_xtuple_0 X0) \Rightarrow (np_1 \in k1_relset_1 k5_numbers X0))$$