

t31_funct_7
(TMK4LKvuVg2fyssHfWEXgVSx2riezQEJoWP)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. X0 \in k9_xtuple_0 (k16_funcop_1 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k1_funct_1 (k16_funcop_1 X0 X1) X0 = X1 \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. \\ & ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 X1) \Rightarrow (k1_funct_1 \\ & (k1_funct_4 X2 X1) X0 = k1_funct_1 X1 X0))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k16_funcop_1 X0 X1)) \wedge (v1_funct_1 (k16_funcop_1 X0 X1)) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & ((X1 \in k9_xtuple_0 X0) \Rightarrow (k2_funct_7 X0 X1 X2 = k1_funct_4 X0 (k16_funcop_1 \\ & X1 X2))) \wedge ((\neg X1 \in k9_xtuple_0 X0) \Rightarrow (k2_funct_7 X0 X1 X2 = X0))) \end{aligned} \quad (5)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & (X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X2 = X1)) \end{aligned}$$