

t31_fuzzy_1

(TMGbkay9NFnQzkn9ie5MJ3UQ9JD4oVjbRZa)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rcomp_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_fuzzy_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_fuzzy_1 : \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v5_relat_1 X1 (k1_rcomp_1 \\ k6_numbers np_1)) \wedge ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k1_numbers) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow \\ (r1_fuzzy_1 (k4_fuzzy_1 X0) X1)) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v5_relat_1 X1 (k1_rcomp_1 \\ k6_numbers np_1)) \wedge ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k1_numbers) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow \\ (\forall X2. ((v5_relat_1 X2 (k1_rcomp_1 k6_numbers np_1)) \wedge (\\ (v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 k1_numbers) \wedge (m1_subset_1 \\ X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow (((r1_fuzzy_1 \\ X1 X2) \wedge (r1_fuzzy_1 X2 X1)) \Rightarrow (r2_funct_2 X0 k1_numbers X1 X2)))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0. (\neg v1_xboole_0 X0) \Rightarrow ((v5_relat_1 (k4_fuzzy_1 X0) (k1_rcomp_1 \\ k6_numbers np_1)) \wedge ((v1_funct_1 (k4_fuzzy_1 X0)) \wedge ((v1_funct_2 \\ (k4_fuzzy_1 X0) X0 k1_numbers) \wedge (m1_subset_1 (k4_fuzzy_1 X0) (\\ k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v5_relat_1 X1 (k1_rcomp_1 \\ k6_numbers np_1)) \wedge ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 X0 k1_numbers) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))))) \Rightarrow \\ ((r1_fuzzy_1 X1 (k4_fuzzy_1 X0)) \Rightarrow (r2_funct_2 X0 k1_numbers X1 \\ (k4_fuzzy_1 X0)))) \end{aligned}$$