

t31\_pdiff\_5  
(TMYqqSDUpUatprBUfoUhccQczctJTpaFs4H)

October 27, 2020

Let  $m2\_finseq\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k1\_euclid : \iota \Rightarrow \iota$  be given. Let  $np\_3 : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r4\_pdiff\_5 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_pdiff\_5 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k11\_pdiff\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k1\_pdiff\_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k11\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_fdiff\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_pdiff\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_rcomp\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_fdiff\_1 : \iota \Rightarrow o$  be given. Let  $v2\_fdiff\_1 : \iota \Rightarrow o$  be given. Let  $k9\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_seq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned}
& \forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 \\
& X1 k1\_numbers) \Rightarrow (\forall X2.(m1\_subset\_1 X2 k1\_numbers) \Rightarrow (\forall X3. \\
& (m2\_finseq\_2 X3 k1\_numbers (k1\_euclid np\_3)) \Rightarrow (\forall X4.(( \\
& v1\_funct\_1 X4) \wedge (m1\_subset\_1 X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k1\_euclid \\
& np\_3) k1\_numbers)))) \Rightarrow ((X3 = k11\_finseq\_1 X0 X1 X2) \Rightarrow (k11\_pdiff\_1 \\
& np\_3 np\_1 X4 X3 = k1\_fdiff\_1 (k1\_pdiff\_2 np\_3 np\_1 X4 X3) X0))))))
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(m1\_subset\_1 X0 k1\_numbers) \Rightarrow (\forall X1.(m1\_subset\_1 \\
& X1 k1\_numbers) \Rightarrow (\forall X2.(m1\_subset\_1 X2 k1\_numbers) \Rightarrow (\forall X3. \\
& (m2\_finseq\_2 X3 k1\_numbers (k1\_euclid np\_3)) \Rightarrow (\forall X4.(( \\
& v1\_funct\_1 X4) \wedge (m1\_subset\_1 X4 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (k1\_euclid \\
& np\_3) k1\_numbers)))) \Rightarrow (((X3 = k11\_finseq\_1 X0 X1 X2) \wedge (r4\_pdiff\_5 \\
& X4 X3)) \Rightarrow (k4\_pdiff\_5 X4 X3 = k1\_fdiff\_1 (k1\_pdiff\_2 np\_3 np\_1 ( \\
& k1\_pdiff\_3 np\_2 np\_3 X4) X3) X0))))))
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & ((v2\_xreal\_0 \ np\_3) \wedge (m2\_subset\_1 \ np\_3 \ k1\_numbers \ k5\_numbers)) \wedge \\ & ((m1\_subset\_1 \ np\_3 \ k5\_numbers) \wedge (m1\_subset\_1 \ np\_3 \ k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\neg v1\_xboole\_0 \ np\_3 \quad (4)$$

Assume the following.

$$\begin{aligned} & ((v2\_xreal\_0 \ np\_2) \wedge (m2\_subset\_1 \ np\_2 \ k1\_numbers \ k5\_numbers)) \wedge \\ & ((m1\_subset\_1 \ np\_2 \ k5\_numbers) \wedge (m1\_subset\_1 \ np\_2 \ k1\_numbers)) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((m1\_subset\_1 \ X0 \ k5\_numbers) \wedge \\ & (((\neg v1\_xboole\_0 \ X1) \wedge (m1\_subset\_1 \ X1 \ k5\_numbers)) \wedge ((v1\_funct\_1 \\ & \ X2) \wedge (m1\_subset\_1 \ X2 \ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \ (k1\_euclid \ X1) \\ & \ k1\_numbers)))))) \Rightarrow ((v1\_funct\_1 \ (k1\_pdiff\_3 \ X0 \ X1 \ X2)) \wedge ((v1\_funct\_2 \\ & \ (k1\_pdiff\_3 \ X0 \ X1 \ X2) \ (k1\_euclid \ X1) \ k1\_numbers) \wedge (m1\_subset\_1 \\ & \ (k1\_pdiff\_3 \ X0 \ X1 \ X2) \ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \ (k1\_euclid \ X1) \\ & \ k1\_numbers)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1\_funct\_1 \ X0) \wedge (m1\_subset\_1 \ X0 \ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \\ & \ (k1\_euclid \ np\_3) \ k1\_numbers)))) \Rightarrow (\forall X1. (m2\_finseq\_2 \ X1 \\ & \ k1\_numbers \ (k1\_euclid \ np\_3)) \Rightarrow ((r4\_pdiff\_5 \ X0 \ X1) \Leftrightarrow (\exists X2. \\ & \ (m1\_subset\_1 \ X2 \ k1\_numbers) \wedge (\exists X3. (m1\_subset\_1 \ X3 \ k1\_numbers) \wedge \\ & \ (\exists X4. (m1\_subset\_1 \ X4 \ k1\_numbers) \wedge ((X1 = k11\_finseq\_1 \ X2 \\ & \ X3 \ X4) \wedge (\exists X5. (m1\_rcomp\_1 \ X5 \ X2) \wedge (r1\_tarski \ X5 \ (k1\_relset\_1 \\ & \ k1\_numbers \ (k1\_pdiff\_2 \ np\_3 \ np\_1 \ (k1\_pdiff\_3 \ np\_2 \ np\_3 \ X0) \\ & \ X1)))) \wedge (\exists X6. ((v1\_funct\_1 \ X6) \wedge (v3\_fdiff\_1 \ X6) \wedge (m1\_subset\_1 \\ & \ X6 \ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \ k1\_numbers \ k1\_numbers)))))) \wedge (\exists X7. \\ & \ ((v1\_funct\_1 \ X7) \wedge ((v2\_fdiff\_1 \ X7) \wedge (m1\_subset\_1 \ X7 \ (k1\_zfmisc\_1 \\ & \ (k2\_zfmisc\_1 \ k1\_numbers \ k1\_numbers)))))) \wedge (\forall X8. (m1\_subset\_1 \\ & \ X8 \ k1\_numbers) \Rightarrow ((X8 \in X5) \Rightarrow (k9\_real\_1 \ (k1\_seq\_1 \ (k1\_pdiff\_2 \ np\_3 \\ & \ np\_1 \ (k1\_pdiff\_3 \ np\_2 \ np\_3 \ X0) \ X1) \ X8) \ (k1\_seq\_1 \ (k1\_pdiff\_2 \\ & \ np\_3 \ np\_1 \ (k1\_pdiff\_3 \ np\_2 \ np\_3 \ X0) \ X1) \ X2) = k7\_real\_1 \ (k1\_seq\_1 \\ & \ X6 \ (k9\_real\_1 \ X8 \ X2)) \ (k1\_seq\_1 \ X7 \ (k9\_real\_1 \ X8 \ X2)))))))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. k11\_finseq\_1 \ X0 \ X1 \ X2 = k7\_finseq\_1 \\ & (k7\_finseq\_1 \ (k9\_finseq\_1 \ X0) \ (k9\_finseq\_1 \ X1)) \ (k9\_finseq\_1 \\ & \ X2) \end{aligned} \quad (8)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(m2\_finseq\_2 X0 k1\_numbers (k1\_euclid np\_3)) \Rightarrow (\forall X1. \\ & ((v1\_funct\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 ( \\ & k1\_euclid np\_3) k1\_numbers)))) \Rightarrow ((r4\_pdf\_5 X1 X0) \Rightarrow (k4\_pdf\_5 \\ & X1 X0 = k11\_pdf\_1 np\_3 np\_1 (k1\_pdf\_3 np\_2 np\_3 X1) X0))) \end{aligned}$$