

t31_rfunct_2 (TMM-
rhkjWym2GHb7tAqEunVQwTkqN7xMieMu)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v8_valued_0 : \iota \Rightarrow o$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\neg(X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(r1_tarski X0 X1) \Rightarrow (r1_tarski (k3_xboole_0 X0 X2) (k3_xboole_0 X1 X2)) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & (k2_zfmisc_1 k1_numbers k1_numbers)))) \Rightarrow ((v8_valued_0 (k2_partfun1 \\ & k1_numbers k1_numbers X1 X0)) \Leftrightarrow (\forall X2. (m1_subset_1 X2 k1_numbers) \Rightarrow \\ & (\forall X3. (m1_subset_1 X3 k1_numbers) \Rightarrow (((X2 \in k9_subset_1 k1_numbers \\ & X0 (k1_relset_1 k1_numbers X1)) \wedge (X3 \in k9_subset_1 k1_numbers X0 \\ & (k1_relset_1 k1_numbers X1))) \Rightarrow ((r1_xreal_0 X3 X2) \vee (r1_xreal_0 \\ & (k1_seq_1 X1 X3) (k1_seq_1 X1 X2))))))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (m1_subset_1 (k1_relset_1 X0 X1) (k1_zfmisc_1 X0)) \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (9)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (10)$$

Theorem 1

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1_funct_1 X2) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers)))) \Rightarrow (((r1_tarski \\ & X0 X1) \wedge (v8_valued_0 (k2_partfun1 k1_numbers k1_numbers X2 X1))) \Rightarrow \\ & (v8_valued_0 (k2_partfun1 k1_numbers k1_numbers X2 X0))) \end{aligned}$$