

# t31\_sheffer2 (TMHXJbMJZZWQiH- hDYMZYzgLkTo3ywWAM2MpH)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v1\_sheffer2 : \iota \Rightarrow o$  be given. Let  $l1\_sheffer1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k5\_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned}
 & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v1\_sheffer2 X0) \wedge (l1\_sheffer1 \\
 & \quad X0))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2. \\
 & \quad (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3.(m1\_subset\_1 X3 \\
 & \quad (u1\_struct\_0 X0)) \Rightarrow (\forall X4.(m1\_subset\_1 X4 (u1\_struct\_0 X0)) \Rightarrow \\
 & \quad (k5\_sheffer1 X0 (k5\_sheffer1 X0 X1 (k5\_sheffer1 X0 X2 X3)) (k5\_sheffer1 \\
 & \quad X0 X1 (k5\_sheffer1 X0 X4 (k5\_sheffer1 X0 X2 X1))) = k5\_sheffer1 X0 \\
 & \quad (k5\_sheffer1 X0 X1 (k5\_sheffer1 X0 X2 X3)) (k5\_sheffer1 X0 X2 X1)))))) \\
 & \hspace{15em} (1)
 \end{aligned}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v1\_sheffer2 X0) \wedge (l1\_sheffer1 \\
 & \quad X0))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2. \\
 & \quad (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3.(m1\_subset\_1 X3 \\
 & \quad (u1\_struct\_0 X0)) \Rightarrow (k5\_sheffer1 X0 (k5\_sheffer1 X0 X1 (k5\_sheffer1 \\
 & \quad X0 X2 X3)) (k5\_sheffer1 X0 X2 X1) = X1)))) \\
 & \hspace{15em} (2)
 \end{aligned}$$

## Theorem 1

$$\begin{aligned}
 & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v1\_sheffer2 X0) \wedge (l1\_sheffer1 \\
 & \quad X0))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 X0)) \Rightarrow (\forall X2. \\
 & \quad (m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3.(m1\_subset\_1 X3 \\
 & \quad (u1\_struct\_0 X0)) \Rightarrow (\forall X4.(m1\_subset\_1 X4 (u1\_struct\_0 X0)) \Rightarrow \\
 & \quad (k5\_sheffer1 X0 (k5\_sheffer1 X0 X1 (k5\_sheffer1 X0 X2 X3)) (k5\_sheffer1 \\
 & \quad X0 X1 (k5\_sheffer1 X0 X4 (k5\_sheffer1 X0 X2 X1))) = X1))))
 \end{aligned}$$