

t32_enumset1
(TMb7TCGTgqXJfoF5xhBAWvxcdnanVLj7FcV)

October 27, 2020

Let $k3_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_enumset1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.k2_enumset1\ X0\ X1\ X2\ X3 = k2_xboole_0\ (k1_enumset1\ X0\ X1\ X2)\ (k1_tarski\ X3) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.k2_enumset1\ X0\ X0\ X1\ X2 = k1_enumset1\ X0\ X1\ X2 \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.k3_enumset1\ X0\ X1\ X2\ X3\ X4 = k2_xboole_0\ (k2_enumset1\ X0\ X1\ X2\ X3)\ (k1_tarski\ X4) \quad (3)$$

Theorem 1

$$\forall X0.\forall X1.\forall X2.\forall X3.k3_enumset1\ X0\ X0\ X1\ X2\ X3 = k2_enumset1\ X0\ X1\ X2\ X3$$