

t32_funct_4

(TMKpS9JMBVhB6LkEs3hhANSjbuZ7GvByrx7)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_xboole_0 (k9_xtuple_0 \\ X0) (k9_xtuple_0 X1)) \Rightarrow (r1_partfun1 X0 X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_partfun1 X0 X1) \Leftrightarrow (r1_tarski \\ X0 (k1_funct_4 X0 X1)))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_xboole_0 (k9_xtuple_0 \\ X0) (k9_xtuple_0 X1)) \Rightarrow (r1_tarski X0 (k1_funct_4 X0 X1)))) \end{aligned}$$