

t32_funct_7

(TMcj2X7wnhZgHFk6cMkvhcJ2m3ATazg8ALu)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ \forall X3. (X1 \neq X3) \Rightarrow (k1_funct_1 (k1_funct_4 X0 (k16_funcop_1 \\ X1 X2)) X3 = k1_funct_1 X0 X3)) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ ((X1 \in k9_xtuple_0 X0) \Rightarrow (k2_funct_7 X0 X1 X2 = k1_funct_4 X0 (k16_funcop_1 \\ X1 X2))) \wedge ((\neg X1 \in k9_xtuple_0 X0) \Rightarrow (k2_funct_7 X0 X1 X2 = X0))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ \forall X3. (X2 \neq X3) \Rightarrow (k1_funct_1 (k2_funct_7 X0 X2 X1) X3 = k1_funct_1 \\ X0 X3)) \end{aligned}$$