

t32_grfunc_1
(TMbX7ir4uSPkU6WN6ScAftgU8JH4SzsgHUr)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg(X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. (v1_relat_1 X1) \Rightarrow (r1_tarski (k6_subset_1 (k9_xtuple_0 X0) (k9_xtuple_0 X1)) (k9_xtuple_0 (k6_subset_1 X0 X1)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (5)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((r1_tarski X0 X1) \Leftrightarrow ((r1_tarski (k9_xtuple_0 X0) (k9_xtuple_0 X1)) \wedge (\forall X2. (X2 \in k9_xtuple_0 X0) \Rightarrow (k1_funct_1 X0 X2 = k1_funct_1 X1 X2)))))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.m1_subset_1 (k6_subset_1 X0 X1) (k1_zfmisc_1 X0) \quad (7)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_funct_1 X1)) \quad (8)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_relat_1 X1)) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall X0.\forall X1.((v1_relat_1 X1)\wedge(v1_funct_1 X1))\Rightarrow(\forall X2. \\ & ((v1_relat_1 X2)\wedge(v1_funct_1 X2))\Rightarrow((X0 \in k6_subset_1 (k9_xtuple_0 \\ & X1) (k9_xtuple_0 X2))\Rightarrow(k1_funct_1 (k6_subset_1 X1 X2) X0 = k1_funct_1 \\ & X1 X0))) \end{aligned}$$