

t32\_helly  
(TMFGpSfSjKtQgAfx8t6snNfJZMEooomrtCJ)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v1\_glib\_000 : \iota \Rightarrow o$  be given. Let  $v3\_glib\_002 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_glib\_000 : \iota \Rightarrow \iota$  be given. Let  $k13\_glib\_001 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_helly : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m3\_glib\_001 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_glib\_001 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_glib\_001 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finset\_1 X0) \wedge (v1\_glib\_000 X0))))) \Rightarrow (\forall X1.(m3\_glib\_001 X1 X0) \Rightarrow (k13\_glib\_001 X0 X1 = k13\_glib\_001 X0 (k6\_glib\_001 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finset\_1 X0) \wedge ((v1\_glib\_000 X0) \wedge (v3\_glib\_002 X0))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k6\_glib\_000 X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (k6\_glib\_000 X0)) \Rightarrow (k6\_glib\_001 X0 (k2\_helly X0 X1 X2) = k2\_helly X0 X2 X1))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finset\_1 X0) \wedge ((v1\_glib\_000 X0) \wedge (v3\_glib\_002 X0))))) \wedge ((m1\_subset\_1 X1 (k6\_glib\_000 X0)) \wedge (m1\_subset\_1 X2 (k6\_glib\_000 X0)))) \Rightarrow ((v5\_glib\_001 (k2\_helly X0 X1 X2) X0) \wedge (m3\_glib\_001 (k2\_helly X0 X1 X2) X0)) \quad (3)$$

**Theorem 1**

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v4\_relat\_1 X0 k5\_numbers) \wedge ((v1\_funct\_1 X0) \wedge ((v1\_finset\_1 X0) \wedge ((v1\_glib\_000 X0) \wedge (v3\_glib\_002 X0))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k6\_glib\_000 X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (k6\_glib\_000 X0)) \Rightarrow (k13\_glib\_001 X0 (k2\_helly X0 X1 X2) = k13\_glib\_001 X0 (k2\_helly X0 X2 X1))))$$