

t32\_nfcont\_3 (TMHbTqQXvGwENJmJRfHfn-  
tXX1b8ZWiYRmHh)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_normsp\_0 : \iota \Rightarrow o$  be given. Let  $v4\_normsp\_0 : \iota \Rightarrow o$  be given. Let  $v2\_normsp\_1 : \iota \Rightarrow o$  be given. Let  $l1\_normsp\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $v1\_nfcont\_3 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_zfmisc\_1 : \iota \Rightarrow o$  be given. Let  $v3\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v5\_relat\_1 X1 X0)) \Rightarrow (k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \quad (1)$$

Assume the following.

$$\forall X0. v1\_zfmisc\_1 (k1\_tarski X0) \quad (2)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (\neg v3\_funct\_1 X0))) \Rightarrow (\neg v1\_zfmisc\_1 (k10\_xtuple\_0 X0)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \quad (5)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\
& X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\
& ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge ((v3\_normsp\_0 \\
& X0) \wedge ((v4\_normsp\_0 X0) \wedge ((v2\_normsp\_1 X0) \wedge (l1\_normsp\_1 X0)))))))))) \Rightarrow \\
& (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k1\_numbers \\
& (u1\_struct\_0 X0)))) \Rightarrow (((v1\_funct\_1 X1) \wedge (v3\_funct\_1 X1)) \Rightarrow ((v1\_funct\_1 \\
& X1) \wedge (v1\_nfcont\_3 X1 X0))))
\end{aligned} \tag{6}$$

**Theorem 1**

$$\begin{aligned}
& \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\
& X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\
& ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge ((v3\_normsp\_0 \\
& X0) \wedge ((v4\_normsp\_0 X0) \wedge ((v2\_normsp\_1 X0) \wedge (l1\_normsp\_1 X0)))))))))) \Rightarrow \\
& (\forall X1.((v1\_funct\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\
& k1\_numbers (u1\_struct\_0 X0)))) \Rightarrow ((\exists X2.(m1\_subset\_1 X2 \\
& (u1\_struct\_0 X0)) \wedge (k2\_relset\_1 (u1\_struct\_0 X0) X1 = k1\_tarski \\
& X2)) \Rightarrow (v1\_nfcont\_3 X1 X0)))
\end{aligned}$$