

t32_quatern3 (TMMR- rxJ5WnhWxtYUeWX9KvCsgLjiQ7oYnDT)

October 27, 2020

Let $v1_quaterni : \iota \Rightarrow o$ be given. Let $k3_quatern2 : \iota \Rightarrow \iota$ be given. Let $k27_quaterni : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k31_quaterni : \iota \Rightarrow \iota$ be given. Let $k32_quaterni : \iota \Rightarrow \iota$ be given. Let $k30_quaterni : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_quaterni : \iota$ be given. Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (k32_quaterni (k31_quaterni X0) = k32_quaterni X0) \quad (1)$$

Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (\forall X1.(v1_quaterni X1) \Rightarrow (k31_quaterni (k27_quaterni X0 X1) = k27_quaterni (k31_quaterni X1) (k31_quaterni X0))) \quad (2)$$

Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (k3_quatern2 X0 = k32_quaterni X0) \quad (3)$$

Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (k31_quaterni X0 = k30_quaterni X0) \quad (4)$$

Assume the following.

$$\forall X0.(v1_quaterni X0) \Rightarrow (v1_quaterni (k30_quaterni X0)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_quaterni X0) \wedge (v1_quaterni X1)) \Rightarrow (m1_subset_1 (k27_quaterni X0 X1) k1_quaterni) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_quaterni) \Rightarrow (v1_quaterni X0) \quad (7)$$

Theorem 1

$$\forall X0.(v1_quaterni X0) \Rightarrow (k3_quatern2 (k27_quaterni X0 X0) = k3_quatern2 (k27_quaterni (k31_quaterni X0) (k31_quaterni X0)))$$