

t32_twoscomp (TMbN8nfin2rAvNfNJSWz1cLeU9PkpUir95U)

October 27, 2020

Let $v1_xtuple_0 : \iota \Rightarrow o$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_twoscomp : \iota$ be given. Let $k3_msafree2 : \iota \Rightarrow \iota$ be given. Let $k37_twoscomp : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k2_msafree2 : \iota \Rightarrow \iota$ be given. Let $k5_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((k2_msafree2 (k5_circcomb X0 X1) = k10_xtuple_0 X1) \wedge (k3_msafree2 (k5_circcomb X0 X1) = k1_tarski (k4_tarski X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k10_finseq_1 X0 X1)) \wedge (v1_funct_1 (k10_finseq_1 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. v1_finseq_1 (k10_finseq_1 X0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k37_twoscomp X0 X1 = k5_circcomb k3_twoscomp (k10_finseq_1 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (5)$$

Theorem 1

$$\forall X0. (\neg v1_xtuple_0 X0) \Rightarrow (\forall X1. (\neg v1_xtuple_0 X1) \Rightarrow (k4_tarski (k10_finseq_1 X0 X1) k3_twoscomp \in k3_msafree2 (k37_twoscomp X0 X1)))$$