

t32_xtuple_0
(TMVYeE3pwWesJhvBc4d3R2sDRY3P4JghEMS)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k11_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((r1_tarski X0 X1) \wedge (r1_tarski X0 X2)) \Rightarrow (r1_tarski X0 (k3_xboole_0 X1 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. r1_tarski (k3_xboole_0 X0 X1) X0 \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (r1_tarski (k11_xtuple_0 X0) (k11_xtuple_0 X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (4)$$

Theorem 1

$$\forall X0. \forall X1. r1_tarski (k11_xtuple_0 (k3_xboole_0 X0 X1)) (k3_xboole_0 (k11_xtuple_0 X0) (k11_xtuple_0 X1))$$