

## t33\_bvfunc25

(TMaxwftTyUvgaev53S4Z2DoigjqDdcMzsLi)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_margrel1 : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_bvfunc\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_bvfunc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge \\ & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\ & (k9\_bvfunc\_1 X0 X2 (k9\_bvfunc\_1 X0 X1 X2)) (k12\_bvfunc\_1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((\neg v1\_xboole\_0 X0) \wedge (((v1\_funct\_1 \\ & X1) \wedge ((v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \wedge ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ & X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 \\ & X0 k6\_margrel1)))))) \Rightarrow ((v1\_funct\_1 (k9\_bvfunc\_1 X0 X1 X2)) \wedge ( \\ & (v1\_funct\_2 (k9\_bvfunc\_1 X0 X1 X2) X0 k6\_margrel1) \wedge (m1\_subset\_1 \\ & (k9\_bvfunc\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \end{aligned} \quad (2)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1.((v1\_funct\_1 X1) \wedge ( \\ & (v1\_funct\_2 X1 X0 k6\_margrel1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 ( \\ & k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge \\ & ((v1\_funct\_2 X2 X0 k6\_margrel1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 k6\_margrel1)))))) \Rightarrow (r2\_funct\_2 X0 k6\_margrel1 \\ & (k9\_bvfunc\_1 X0 X1 (k9\_bvfunc\_1 X0 (k9\_bvfunc\_1 X0 X1 X2) X1)) (k12\_bvfunc\_1 \\ & X0))) \end{aligned}$$