

t33_bvfunc_5

(TMd6EhNSQhonYPYL5RdkcB6d3NBuVuimcVB)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_bvfunc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_bvfunc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\ & (k9_bvfunc_1 X0 (k9_bvfunc_1 X0 (k1_bvfunc_1 X0 X1) (k1_bvfunc_1 \\ & X0 X2)) (k9_bvfunc_1 X0 X2 X1)) (k12_bvfunc_1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (k1_bvfunc_1 X0 (k1_bvfunc_1 \\ & X0 X1) = X1) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((v1_funct_1 X1) \wedge (\\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow ((v1_funct_1 (k1_bvfunc_1 \\ & X0 X1)) \wedge ((v1_funct_2 (k1_bvfunc_1 X0 X1) X0 k6_margrel1) \wedge (m1_subset_1 \\ & (k1_bvfunc_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k6_margrel1)))))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge \\ & (v1_funct_2 X1 X0 k6_margrel1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (\\ & k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge \\ & ((v1_funct_2 X2 X0 k6_margrel1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 k6_margrel1)))))) \Rightarrow (r2_funct_2 X0 k6_margrel1 \\ & (k9_bvfunc_1 X0 (k9_bvfunc_1 X0 X1 (k1_bvfunc_1 X0 X2)) (k9_bvfunc_1 \\ & X0 X2 (k1_bvfunc_1 X0 X1)) (k12_bvfunc_1 X0)))) \end{aligned}$$