

t33_cqc_the3

(TMbD7TASmcWrbDmYFGefzbhUhF8A82Snx5r)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_cqc_lang : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $r3_cqc_the1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_cqc_lang : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow ((r3_cqc_the1 X0 X3 (k7_cqc_lang X0 X1 X2)) \Rightarrow \\ (r3_cqc_the1 X0 X3 X1) \wedge (r3_cqc_the1 X0 X3 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow (((r3_cqc_the1 X0 X3 X1) \wedge (r3_cqc_the1 X0 X3 \\ X2)) \Rightarrow (r3_cqc_the1 X0 X3 (k7_cqc_lang X0 X1 X2)))))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k9_qc_lang1 \\ X0) (k3_cqc_lang X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 \\ (k3_cqc_lang X0))) \Rightarrow ((r3_cqc_the1 X0 X3 (k7_cqc_lang X0 X1 X2)) \Leftrightarrow \\ (r3_cqc_the1 X0 X3 X1) \wedge (r3_cqc_the1 X0 X3 X2)))))) \end{aligned}$$