

t33_finseq_5

(TMLvuFS9Nypiius6a3nCykkPMnT2xWVES4p)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_rfinseq : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_rfinseq : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v7_ordinal1 X1) \wedge (m1_finseq_1 X2 X0)) \Rightarrow (k2_rfinseq X0 X1 X2 = k1_rfinseq X1 X2) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (v1_funct_1 X1) \wedge (v1_finseq_1 X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((v7_ordinal1 X1) \wedge (m1_finseq_1 X2 X0)) \Rightarrow (m2_finseq_1 (k2_rfinseq X0 X1 X2) X0) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((m1_finseq_1 X1 X0) \Leftrightarrow (r1_tarski (k10_xtuple_0 X1) X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarSKI X0 X1)\Leftrightarrow(\forall X2.(X2 \in X0)\Rightarrow (X2 \in X1)) \quad (7)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow(\forall X2.(m2_finseq_1 X2 X1)\Rightarrow(r1_tarSKI (k10_xtuple_0 (k2_rfinseq X1 X0 X2)) (k10_xtuple_0 X2))))$$