

t33_funct_8

(TMXy4muddgZy5HgeqMFyatNdx7yqzB6nB2z)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v6_funct_8 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k32_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k30_valued_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_funct_8 : \iota \Rightarrow o$ be given. Let $v1_funct_8 : \iota \Rightarrow o$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $v5_funct_8 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_real_1 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ & ((k9_xtuple_0 (k30_valued_1 X0) = k9_xtuple_0 X0) \wedge (\forall X1. \\ & k1_funct_1 (k30_valued_1 X0) X1 = k4_xcmplx_0 (k1_funct_1 X0 X1))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v3_membered X1) \wedge ((v1_funct_1 \\ & X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow (k32_valued_1 \\ & X0 X1 X2 = k30_valued_1 X2) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (\\ & k1_relset_1 X0 X1 = k9_xtuple_0 X1) \end{aligned} \tag{3}$$

Assume the following.

$$v3_membered k1_numbers \tag{4}$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge (v2_funct_8 X0)) \Rightarrow (v1_funct_8 (k9_xtuple_0 \\ & X0)) \end{aligned} \tag{5}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((v3_membered\ X1)\wedge((v1_funct_1 \\ X2)\wedge(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1))))\Rightarrow((v1_funct_1 \\ (k32_valued_1\ X0\ X1\ X2))\wedge(m1_subset_1\ (k32_valued_1\ X0\ X1\ X2)\ (\\ k1_zfmisc_1\ (k2_zfmisc_1\ X0\ k1_numbers)))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_valued_0\ X0)))\Rightarrow \\ ((v1_relat_1\ (k30_valued_1\ X0))\wedge((v1_funct_1\ (k30_valued_1\ \\ X0))\wedge(v1_valued_0\ (k30_valued_1\ X0)))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_membered\ X0)\Rightarrow(\forall X1.(v1_membered\ X1)\Rightarrow(\forall X2. \\ ((v1_funct_1\ X2)\wedge(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0 \\ X1))))\Rightarrow((v5_funct_8\ X2\ X0\ X1)\Leftrightarrow(\forall X3.(m1_subset_1\ X3\ k1_numbers)\Rightarrow \\ (((X3\in k1_relset_1\ X0\ X2)\wedge(k1_real_1\ X3\in k1_relset_1\ X0\ X2))\Rightarrow(\\ k1_funct_1\ X2\ (k1_real_1\ X3) = k4_xcmplx_0\ (k1_funct_1\ X2\ X3)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.(v1_relat_1\ X0)\Rightarrow((v2_funct_8\ X0)\Leftrightarrow(v1_funct_8\ (k9_xtuple_0 \\ X0))) \quad (9)$$

Assume the following.

$$\forall X0.(v3_membered\ X0)\Rightarrow(v1_membered\ X0) \quad (10)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1_membered\ X0)\wedge(v1_membered\ X1))\Rightarrow(\\ \forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow \\ (((v1_funct_1\ X2)\wedge(v6_funct_8\ X2\ X0\ X1))\Rightarrow((v1_funct_1\ X2)\wedge((\\ v2_funct_8\ X2)\wedge(v5_funct_8\ X2\ X0\ X1)))) \end{aligned} \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((v1_membered\ X0)\wedge(v1_membered\ X1))\Rightarrow(\\ \forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow \\ (((v1_funct_1\ X2)\wedge((v2_funct_8\ X2)\wedge(v5_funct_8\ X2\ X0\ X1)))\Rightarrow(\\ (v1_funct_1\ X2)\wedge(v6_funct_8\ X2\ X0\ X1)))) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1 \\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow((v4_relat_1\ X2\ X0)\wedge(v5_relat_1\ X2\ X1)) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.(v1_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_valued_0 X2)) \quad (15)$$

Theorem 1

$$\forall X0.((v1_funct_1 X0)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers k1_numbers))))\Rightarrow((v6_funct_8 X0 k1_numbers k1_numbers)\Rightarrow(v6_funct_8 (k32_valued_1 k1_numbers k1_numbers X0) k1_numbers k1_numbers))$$