

# t33\_jgraph\_7 (TMGNRdvFsAwtPCb- JjM6dkQF9vSTmn8aD7kk)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k17\_euclid : \iota \Rightarrow \iota$  be given. Let  $k18\_euclid : \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_jordan17 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_sppol\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_jordan6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Assume the following.

$$\begin{aligned}
& \forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 (k15\_euclid np\_2))) \Rightarrow \\
& (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 (k15\_euclid np\_2))) \Rightarrow \\
& (\forall X2.(v1\_xreal\_0 X2) \Rightarrow (\forall X3.(v1\_xreal\_0 X3) \Rightarrow ((( \\
& r1\_xxreal\_0 (k18\_euclid X1) (k18\_euclid X0)) \wedge ((r1\_xxreal\_0 ( \\
& k18\_euclid X0) X3) \wedge (r1\_xxreal\_0 (k17\_euclid X1) X2))) \Rightarrow ((r1\_xxreal\_0 \\
& X3 (k18\_euclid X1)) \vee ((r1\_xxreal\_0 (k17\_euclid X1) (k17\_euclid \\
& X0)) \vee (r1\_jordan6 (k1\_sppol\_2 (k17\_euclid X0) X2 (k18\_euclid X1) \\
& X3) X0 X1))))))
\end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\
& (v1\_xxreal\_0 X2) \Rightarrow (((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow \\
& (r1\_xxreal\_0 X0 X2))))
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 (k15\_euclid np\_2))) \Rightarrow \\
& (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 (k15\_euclid np\_2))) \Rightarrow \\
& (\forall X2.(v1\_xreal\_0 X2) \Rightarrow (\forall X3.(v1\_xreal\_0 X3) \Rightarrow (\forall X4. \\
& (v1\_xreal\_0 X4) \Rightarrow (\forall X5.(v1\_xreal\_0 X5) \Rightarrow (((k18\_euclid X0 = \\
& X4) \wedge ((k18\_euclid X1 = X4) \wedge (r1\_xxreal\_0 (k17\_euclid X0) X3))) \Rightarrow \\
& ((r1\_xxreal\_0 X3 X2) \vee ((r1\_xxreal\_0 X5 X4) \vee ((r1\_xxreal\_0 (k17\_euclid \\
& X1) X2) \vee ((r1\_xxreal\_0 (k17\_euclid X0) (k17\_euclid X1)) \vee (r1\_jordan6 \\
& (k1\_sppol\_2 X2 X3 X4 X5) X0 X1))))))))))
\end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1\_xreal\_0 X0)\wedge \\ & ((v1\_xreal\_0 X1)\wedge((v1\_xreal\_0 X2)\wedge(v1\_xreal\_0 X3))))\Rightarrow(m1\_subset\_1 \\ & (k1\_sppol\_2 X0 X1 X2 X3) (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid \\ & np\_2)))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 (k15\_euclid np\_2)))\Rightarrow \\ & (m1\_subset\_1 (k17\_euclid X0) k1\_numbers) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid \\ & np\_2))))\Rightarrow(\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 (k15\_euclid \\ & np\_2))))\Rightarrow(\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 (k15\_euclid \\ & np\_2))))\Rightarrow(\forall X3.(m1\_subset\_1 X3 (u1\_struct\_0 (k15\_euclid \\ & np\_2))))\Rightarrow(\forall X4.(m1\_subset\_1 X4 (u1\_struct\_0 (k15\_euclid \\ & np\_2))))\Rightarrow((r1\_jordan17 X0 X1 X2 X3 X4)\Leftrightarrow(\neg(\neg(r1\_jordan6 X0 X1 X2)\wedge \\ & ((r1\_jordan6 X0 X2 X3)\wedge(r1\_jordan6 X0 X3 X4))\wedge(\neg(r1\_jordan6 \\ & X0 X2 X3)\wedge(r1\_jordan6 X0 X3 X4)\wedge(r1\_jordan6 X0 X4 X1))\wedge(\neg(r1\_jordan6 \\ & X0 X3 X4)\wedge(r1\_jordan6 X0 X4 X1)\wedge(r1\_jordan6 X0 X1 X2))\wedge(\neg(r1\_jordan6 \\ & X0 X4 X1)\wedge((r1\_jordan6 X0 X1 X2)\wedge(r1\_jordan6 X0 X2 X3)))))))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1\_xxreal\_0 X0)\wedge(v1\_xxreal\_0 X1))\Rightarrow( \\ & (r1\_xxreal\_0 X0 X1)\vee(r1\_xxreal\_0 X1 X0)) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow(v1\_xxreal\_0 X0) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (9)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 (k15\_euclid np\_2)))\Rightarrow \\ & (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 (k15\_euclid np\_2))))\Rightarrow \\ & (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 (k15\_euclid np\_2))))\Rightarrow \\ & (\forall X3.(m1\_subset\_1 X3 (u1\_struct\_0 (k15\_euclid np\_2))))\Rightarrow \\ & (\forall X4.(v1\_xreal\_0 X4)\Rightarrow(\forall X5.(v1\_xreal\_0 X5)\Rightarrow(\forall X6. \\ & (v1\_xreal\_0 X6)\Rightarrow(\forall X7.(v1\_xreal\_0 X7)\Rightarrow(((k17\_euclid X0 = \\ & X4)\wedge((k18\_euclid X1 = X6)\wedge((k18\_euclid X2 = X6)\wedge((k18\_euclid X3 = \\ & X6)\wedge((r1\_xxreal\_0 X6 (k18\_euclid X0))\wedge((r1\_xxreal\_0 (k18\_euclid \\ & X0) X7)\wedge(r1\_xxreal\_0 (k17\_euclid X1) X5))))))\Rightarrow((r1\_xxreal\_0 \\ & X5 X4)\vee((r1\_xxreal\_0 X7 X6)\vee((r1\_xxreal\_0 (k17\_euclid X3) X4)\vee \\ & ((r1\_xxreal\_0 (k17\_euclid X2) (k17\_euclid X3))\vee((r1\_xxreal\_0 \\ & (k17\_euclid X1) (k17\_euclid X2))\vee(r1\_jordan17 (k1\_sppol\_2 X4 \\ & X5 X6 X7) X0 X1 X2 X3)))))))))) \end{aligned}$$