

## t33\_jordan21

(TMaigr5ogz6wWQ2DGMKVcmimv9BXoKZ39ea)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $v2\_sppol\_1 : \iota \Rightarrow o$  be given. Let  $k1\_rltopsp1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_jordan21 : \iota \Rightarrow \iota$  be given. Let  $k19\_euclid : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_real\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_pscomp\_1 : \iota \Rightarrow \iota$  be given. Let  $k8\_pscomp\_1 : \iota \Rightarrow \iota$  be given. Let  $k9\_pscomp\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k17\_euclid : \iota \Rightarrow \iota$  be given. Let  $k18\_euclid : \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_seq\_4 : \iota \Rightarrow \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k5\_pscomp\_1 : \iota$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_jordan6 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow ((k17\_euclid (k19\_euclid X0 X1) = X0) \wedge (k18\_euclid (k19\_euclid X0 X1) = X1))) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 (k15\_euclid np\_2))) \Rightarrow \\ & (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 (k15\_euclid np\_2))) \Rightarrow \\ & ((k17\_euclid X0 = k17\_euclid X1) \Leftrightarrow (v2\_sppol\_1 (k1\_rltopsp1 (k15\_euclid \\ & \quad np\_2) X0 X1)))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid np\_2)))) \Rightarrow (m1\_subset\_1 (k9\_pscomp\_1 X0) k1\_numbers) \quad (4)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid np\_2)))) \Rightarrow (m1\_subset\_1 (k8\_pscomp\_1 X0) k1\_numbers) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (m1\_subset\_1 (k7\_relset\_1 X0 X1 X2 X3) (k1\_zfmisc\_1 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers) \wedge (v1\_xreal\_0 X1)) \Rightarrow (m1\_subset\_1 (k7\_real\_1 X0 X1) k1\_numbers) \quad (7)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid np\_2)))) \Rightarrow (m1\_subset\_1 (k6\_pscomp\_1 X0) k1\_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 k1\_numbers)) \Rightarrow (m1\_subset\_1 (k5\_seq\_4 X0) k1\_numbers) \quad (9)$$

Assume the following.

$$(v1\_funct\_1 k5\_pscomp\_1) \wedge ((v1\_funct\_2 k5\_pscomp\_1 (u1\_struct\_0 (k15\_euclid np\_2)) k1\_numbers) \wedge (m1\_subset\_1 k5\_pscomp\_1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid np\_2)) k1\_numbers)))) \quad (10)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid np\_2)))) \Rightarrow (m1\_subset\_1 (k2\_jordan21 X0) (u1\_struct\_0 (k15\_euclid np\_2))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (m1\_subset\_1 (k19\_euclid X0 X1) (u1\_struct\_0 (k15\_euclid np\_2))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers) \wedge (v1\_xreal\_0 X1)) \Rightarrow (m1\_subset\_1 (k10\_real\_1 X0 X1) k1\_numbers) \quad (13)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid np\_2)))) \Rightarrow (k2\_jordan21 X0 = k19\_euclid (k10\_real\_1 (k7\_real\_1 (k8\_pscomp\_1 X0) (k6\_pscomp\_1 X0)) np\_2) (k5\_seq\_4 (k7\_relset\_1 (u1\_struct\_0 (k15\_euclid np\_2)) k1\_numbers k5\_pscomp\_1 (k9\_subset\_1 (u1\_struct\_0 (k15\_euclid np\_2)) X0 (k6\_jordan6 (k10\_real\_1 (k7\_real\_1 (k8\_pscomp\_1 X0) (k6\_pscomp\_1 X0)) np\_2)))))) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_numbers)\wedge(v1\_xreal\_0 X1))\Rightarrow(k7\_real\_1 X0 X1 = k7\_real\_1 X1 X0) \quad (15)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (16)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 (u1\_struct\_0 (k15\_euclid np\_2))))\Rightarrow(v2\_sppol\_1 (k1\_rttopsp1 (k15\_euclid np\_2) (k2\_jordan21 X0) (k19\_euclid (k10\_real\_1 (k7\_real\_1 (k6\_pscomp\_1 X0) (k8\_pscomp\_1 X0)) np\_2) (k9\_pscomp\_1 X0))))$$