

# t33\_matrprob (TMJPAmbisE- Zou9QkgN9R9qLx4GCzKzGSTZh)

October 27, 2020

Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_seq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k14\_rvsum\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v3\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v3\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k11\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v3\_membered : \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (\neg(\neg r1\_xxreal\_0 X0 X1) \wedge ((\neg v3\_xxreal\_0 X1) \wedge (\neg v2\_xxreal\_0 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge ((v3\_valued\_0 X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow (\forall X2.((v1\_relat\_1 X2) \wedge ((v1\_funct\_1 X2) \wedge ((v3\_valued\_0 X2) \wedge (v1\_finseq\_1 X2)))) \Rightarrow (k1\_seq\_1 (k14\_rvsum\_1 X1 X2) X0 = k11\_binop\_2 (k1\_seq\_1 X1 X0) (k1\_seq\_1 X2 X0))) \quad (2)$$

Assume the following.

$$m1\_subset\_1 k1\_xboole\_0 k4\_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.(v1\_xreal\_0 X1) \Rightarrow (((r1\_xxreal\_0 k6\_numbers X0) \wedge (r1\_xxreal\_0 k6\_numbers X1)) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k3\_xcmplx\_0 X0 X1)))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Leftrightarrow(m1\_finseq\_1 X1 X0) \quad (5)$$

Assume the following.

$$k6\_numbers = k1\_xboole\_0 \quad (6)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v3\_valued\_0 X0)))\Rightarrow(k1\_seq\_1 X0 X1 = k1\_funct\_1 X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(k11\_binop\_2 X0 X1 = k3\_xcmplx\_0 X0 X1) \quad (10)$$

Assume the following.

$$v3\_membered k1\_numbers \quad (11)$$

Assume the following.

$$v1\_xboole\_0 k1\_xboole\_0 \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge((v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))) \quad (14)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v3\_valued\_0 X0)))\Rightarrow(m1\_subset\_1 (k1\_seq\_1 X0 X1) k1\_numbers) \quad (16)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ ((X1 \in k9\_xtuple\_0 X0) \Rightarrow ((X2 = k1\_funct\_1 X0 X1) \Leftrightarrow (k4\_tarSKI X1 X2 \in \\ X0))) \wedge ((\neg X1 \in k9\_xtuple\_0 X0) \Rightarrow ((X2 = k1\_funct\_1 X0 X1) \Leftrightarrow (X2 = k1\_xboole\_0)))) \end{aligned} \quad (17)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (18)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 (k1\_zfmisc\_1 k1\_numbers)) \Rightarrow (v3\_membered X0) \quad (19)$$

Assume the following.

$$\forall X0.((v1\_xxreal\_0 X0) \wedge (v2\_xxreal\_0 X0)) \Rightarrow ((\neg v1\_xboole\_0 X0) \wedge ((v1\_xxreal\_0 X0) \wedge (\neg v3\_xxreal\_0 X0))) \quad (20)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((v7\_ordinal1 X0) \wedge (\neg v3\_xxreal\_0 X0)) \quad (21)$$

Assume the following.

$$\forall X0. \forall X1.(m1\_finseq\_1 X1 X0) \Rightarrow (v5\_relat\_1 X1 X0) \quad (22)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v5\_relat\_1 X0 k1\_numbers)) \Rightarrow ((v1\_relat\_1 X0) \wedge (v3\_valued\_0 X0)) \quad (23)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v1\_xxreal\_0 X0) \quad (24)$$

Assume the following.

$$\forall X0.(v3\_membered X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 X0) \Rightarrow (v1\_xreal\_0 X1)) \quad (25)$$

### Theorem 1

$$\begin{aligned} \forall X0.(m2\_finseq\_1 X0 k1\_numbers) \Rightarrow (\forall X1.(m2\_finseq\_1 \\ X1 k1\_numbers) \Rightarrow (((\forall X2.(m2\_subset\_1 X2 k1\_numbers k5\_numbers) \Rightarrow \\ ((X2 \in k4\_finseq\_1 X0) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k1\_seq\_1 X0 X2)))) \wedge \\ (\forall X2.(m2\_subset\_1 X2 k1\_numbers k5\_numbers) \Rightarrow ((X2 \in k4\_finseq\_1 \\ X1) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k1\_seq\_1 X1 X2)))))) \Rightarrow (\forall X2. \\ (m2\_subset\_1 X2 k1\_numbers k5\_numbers) \Rightarrow ((X2 \in k4\_finseq\_1 (k14\_rvsum\_1 \\ X0 X1)) \Rightarrow (r1\_xxreal\_0 k6\_numbers (k1\_seq\_1 (k14\_rvsum\_1 X0 X1) \\ X2)))))) \end{aligned}$$