

t33_nfcont_1

(TMaFUx32X795Yz6LvF_xosaLzBYisphVeqWB)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_normsp_0 : \iota \Rightarrow o$ be given. Let $v4_normsp_0 : \iota \Rightarrow o$ be given. Let $v2_normsp_1 : \iota \Rightarrow o$ be given. Let $l1_normsp_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_nfcont_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r4_nfcont_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_seq_4 : \iota \Rightarrow \iota$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_seq_4 : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_rcomp_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow ((k9_xtuple_0 X0 = k1_xboole_0) \Leftrightarrow (k10_xtuple_0 X0 = k1_xboole_0)) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X3) \wedge \\ & (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \Rightarrow (\neg(X2 \in k10_xtuple_0 \\ & X3) \wedge (\forall X4.(m1_subset_1 X4 X0) \Rightarrow (\neg(X4 \in k1_relset_1 X0 X3) \wedge \\ & (X2 = k1_funct_1 X3 X4)))) \quad (2) \end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\
& X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\
& ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge ((v3_normsp_0 \\
& X0) \wedge ((v4_normsp_0 X0) \wedge ((v2_normsp_1 X0) \wedge (l1_normsp_1 X0)))))))))) \Rightarrow \\
& (\forall X1.((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\
& (u1_struct_0 X0) k1_numbers)))) \Rightarrow (((v1_nfcont_1 (k1_relset_1 \\
& (u1_struct_0 X0) X1) X0) \wedge (r4_nfcont_1 X0 X1 (k1_relset_1 (u1_struct_0 \\
& X0) X1))) \Rightarrow (v1_rcomp_1 (k2_relset_1 k1_numbers X1))))
\end{aligned} \tag{3}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 k1_numbers)) \Rightarrow ((v1_rcomp_1 X0) \Rightarrow ((X0 = k1_xboole_0) \vee ((k4_seq_4 X0 \in X0) \wedge (k5_seq_4 X0 \in X0)))) \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (k2_relset_1 X0 X1 = k10_xtuple_0 X1) \tag{5}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \tag{6}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v5_relat_1 X1 X0)) \Rightarrow (m1_subset_1 (k2_relset_1 X0 X1) (k1_zfmisc_1 X0)) \tag{7}$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge ((v5_relat_1 X1 X0) \wedge (v1_funct_1 X1))) \Rightarrow (\forall X2.(X2 \in k9_xtuple_0 X1) \Rightarrow (k7_partfun1 X0 X1 X2 = k1_funct_1 X1 X2)) \tag{8}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \tag{9}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \tag{10}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\ & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge ((v3_normsp_0 \\ & X0) \wedge ((v4_normsp_0 X0) \wedge ((v2_normsp_1 X0) \wedge (l1_normsp_1 X0)))))))))) \Rightarrow \\ & (\forall X1.((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & (u1_struct_0 X0) k1_numbers)))) \Rightarrow (\neg(k1_relset_1 (u1_struct_0 \\ & X0) X1 \neq k1_xboole_0) \wedge ((v1_nfcont_1 (k1_relset_1 (u1_struct_0 \\ & X0) X1) X0) \wedge ((r4_nfcont_1 X0 X1 (k1_relset_1 (u1_struct_0 X0) X1)) \wedge \\ & (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 \\ & X3 (u1_struct_0 X0)) \Rightarrow (\neg(X2 \in k1_relset_1 (u1_struct_0 X0) X1) \wedge \\ & ((X3 \in k1_relset_1 (u1_struct_0 X0) X1) \wedge ((k7_partfun1 k1_numbers \\ & X1 X2 = k4_seq_4 (k2_relset_1 k1_numbers X1)) \wedge (k7_partfun1 k1_numbers \\ & X1 X3 = k5_seq_4 (k2_relset_1 k1_numbers X1)))))))))) \end{aligned}$$