

## t33\_rusub\_5

(TMSC7mDE4y1uE7HapTx2BuDi1dnGgvGvNcf)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v2\_bhsp\_1 : \iota \Rightarrow o$  be given. Let  $l1\_bhsp\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_bhsp\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\ & X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\ & ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge ((v2\_bhsp\_1 \\ & X0) \wedge (l1\_bhsp\_1 X0)))))))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 \\ & X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (u1\_struct\_0 X0)) \Rightarrow (\forall X3. \\ & (m1\_subset\_1 X3 k1\_numbers) \Rightarrow (\forall X4.(m1\_subset\_1 X4 k1\_numbers) \Rightarrow \\ & (((X1 \in k4\_bhsp\_2 X0 X2 X3) \wedge (r1\_xxreal\_0 X3 X4)) \Rightarrow (X1 \in k4\_bhsp\_2 \\ & X0 X2 X4)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)) \Rightarrow (\forall X2. \\ & (m1\_subset\_1 X2 (k1\_zfmisc\_1 X0)) \Rightarrow ((\forall X3.(m1\_subset\_1 \\ & X3 X0) \Rightarrow ((X3 \in X1) \Rightarrow (X3 \in X2))) \Rightarrow (r1\_tarski X1 X2))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 \\ & X0) \wedge ((v2\_rlvect\_1 X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge \\ & ((v5\_rlvect\_1 X0) \wedge ((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 \\ & X0) \wedge ((v2\_bhsp\_1 X0) \wedge (l1\_bhsp\_1 X0)))))))))) \wedge ((m1\_subset\_1 \\ & X1 (u1\_struct\_0 X0)) \wedge (m1\_subset\_1 X2 k1\_numbers))) \Rightarrow (m1\_subset\_1 \\ & (k4\_bhsp\_2 X0 X1 X2) (k1\_zfmisc\_1 (u1\_struct\_0 X0))) \end{aligned} \tag{3}$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge (v13\_algstr\_0 X0) \wedge (v2\_rlvect\_1 \\ & X0) \wedge (v3\_rlvect\_1 X0) \wedge (v4\_rlvect\_1 X0) \wedge (v5\_rlvect\_1 X0) \wedge \\ & ((v6\_rlvect\_1 X0) \wedge (v7\_rlvect\_1 X0) \wedge (v8\_rlvect\_1 X0) \wedge ((v2\_bhsp\_1 \\ & X0) \wedge (l1\_bhsp\_1 X0)))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (u1\_struct\_0 \\ & X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 k1\_numbers) \Rightarrow (\forall X3.(m1\_subset\_1 \\ & X3 k1\_numbers) \Rightarrow ((r1\_xxreal\_0 X2 X3) \Rightarrow (r1\_tarski (k4\_bhsp\_2 X0 \\ & X1 X2) (k4\_bhsp\_2 X0 X1 X3)))))) \end{aligned}$$